

## WEST Search History

DATE: Wednesday, June 01, 2005

Hide?	<u>Set</u> <u>Name</u>	<u>Query</u>	<u>Hit</u> <u>Count</u>
		<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L1	((peg or pegalate or pegylate or pegylation or pagylating or pegged or pegconjugate or peg\$ or polyethyleneglycol or ethyleneglycol or ethylene-glycol or poly-ethylene-glycol or polyol or polyene or (polymer near5 glycol)).ti,ab,clm. same polysorbate.ti,ab,clm. or poly-sorbate.ti,ab,clm. or \$orbate.ti,ab,clm. same (supposit\$ or composition or vaccine or insertion or application or applying).ti,ab,clm.)	160
<input type="checkbox"/>	L2	l1 and suppos\$	14

END OF SEARCH HISTORY

## Dorlands Medical Dictionary on-line

**Tween (Tween)** (tw[emacr]n) trademark for preparations of polysorbates, used with a numerical suffix; e.g., *Tween 80* is a trademark for polysorbate 80.

**polysorbate (poly·sor·bate)** (pol"e-sor'b[amacr]t) a generic name for esters of sorbitol and its anhydrides condensed with polymers of ethylene oxide, used as surfactant agents: *p. 20*,  $C_{38}H_{114}O_{26}$ , is polyoxyethylene 20 sorbitan monolaurate; *p. 40*,  $C_{62}H_{122}O_{26}$ , is polyoxyethylene 20 sorbitan monopalmitate; *p. 60*,  $C_{64}H_{126}O_{26}$ , is polyethylene 20 sorbitan monostearate; *p. 65*,  $C_{100}H_{194}O_{28}$ , is polyethylene 20 sorbitan tristearate; *p. 80*,  $C_{64}H_{124}O_{26}$ , is polyethylene 20 sorbitan monooleate; *p. 85*,  $C_{100}H_{188}O_{28}$ , is polyethylene 20 sorbitan trioleate. *Polysorbates 20, 40, 60, and 80* are official in NF.

101. 5968490. 11 Sep 98; 19 Oct 99. Antiperspirant deodorant compositions. Sun; Wei Mei, et al. 424/65; 424/400 424/401 424/66 424/68 424/70.1 424/70.21 424/70.27 424/70.28. A61K007/32 A61K007/11 A61K007/00.

---

☐ 102. 5961997. 24 Mar 98; 05 Oct 99. Antipruritic composition. Swinehart; James M.. 424/401; 514/846. A61K007/48.

---

☐ 103. 5922754. 02 Oct 98; 13 Jul 99. Pharmaceutical compositions containing paclitaxel. Burchett; Mark K., et al. 514/449; A61K031/335.

---

☐ 104. 5888492. 21 May 97; 30 Mar 99. Rinse-off skin conditioner. Starch; Mike. 424/78.03; 424/70.19 424/70.21 424/70.31 510/159. A61K031/74 A61K007/50.

---

☐ 105. 5858401. 22 Jan 97; 12 Jan 99. Pharmaceutical composition for cyclosporines. Bhalani; Vinayak T., et al. 424/450; 424/405 514/773. A61K009/48.

---

☐ 106. 5733572. 29 Nov 94; 31 Mar 98. Gas and gaseous precursor filled microspheres as topical and subcutaneous delivery vehicles. Unger; Evan C., et al. 424/450; 424/1.21 424/489 424/9.321 424/9.4 436/829. A61K009/127.

---

☐ 107. 5721275. 21 Jan 92; 24 Feb 98. Slow release vehicles for minimizing skin irritancy of topical compositions. Bazzano; Gail S.. 514/559; 424/78.02 514/859 514/944. A61K031/20 A61K031/78.

---

☐ 108. 5705187. 05 Apr 95; 06 Jan 98. Compositions of lipids and stabilizing materials. Unger; Evan C.. 424/450; 424/43 424/45 424/9.321 424/9.4 428/402.2. A61K009/127.

---

☐ 109. 5681552. 23 May 95; 28 Oct 97. Clear cosmetic stick composition containing a combination of anionic and non-ionic surfactants. Shevade; Makarand, et al. 424/65; 424/78.02 424/78.08 424/78.18 512/1 514/944 514/946. A61K007/32.

---

☐ 110. 5679360. 19 Dec 95; 21 Oct 97. Substance P antagonist for the treatment of lichens, prurigo, pruritus. de Lacharriere; Olivier, et al. 424/401; 424/434 424/436 424/450 424/489 514/2 514/844 514/937 514/944. A61K007/48.

---

☐ 111. 5676956. 10 Apr 96; 14 Oct 97. Composition and method for reducing under-eye puffiness. Duffy; John A., et al. 424/401; 424/195.16 424/520. A61K007/00.

---

☐ 112. 5672340. 07 Mar 96; 30 Sep 97. Antiperspirant deodorant compositions. Sun; Wei Mei, et al. 424/66; 424/65 424/68. A61K007/38 A61K007/34.

---

☐ 113. 5643587. 15 Feb 96; 01 Jul 97. Composition and method for under-eye skin lightening. Scancarella; Neil D., et al. 424/401; 424/195.16 424/62. A61K007/00.

---

☐ 114. 5635166. 22 Apr 96; 03 Jun 97. Antiperspirant deodorant compositions. Galleguillos; Ramiro, et al. 424/66; 424/68. A61K007/34 A61K007/38.

---

☐ 115. 5632974. 06 Jun 96; 27 May 97. Antiperspirant deodorant compositions. Galleguillos; Ramiro, et al. 424/66; 424/68. A61K007/34 A61K007/38.

---

☐ 116. 5591425. 23 Mar 95; 07 Jan 97. Two-package pretreatment and hair relaxer compositions.

---

Dhaliwal; Tehsel. 424/70.4; 424/70.16 424/70.2. A61K007/06 A61K007/09.

---

☐ 117. 5549887. 14 Dec 94; 27 Aug 96. Antiperspirant deodorant compositions. Galleguillos; Ramiro, et al. 424/66; 424/68. A61K007/34 A61K007/38.

---

☐ 118. 5534245. 17 Jan 95; 09 Jul 96. Antiperspirant deodorant compositions. Galleguillos; Ramiro, et al. 424/66; 424/68. A61K007/34 A61K007/38.

---

☐ 119. 5356627. 01 Feb 93; 18 Oct 94. Waterproof cosmetic compositions. Da Cunha; Kathleen, et al. 424/401; 424/63 514/937 514/938. A61K007/02 A61K007/032.

---

☐ 120. 5304370. 09 Jun 92; 19 Apr 94. Hair relaxer composition and associated methods. Hawkins; Geoffrey R., et al. 424/70.2; 132/205 424/70.19 424/70.31. A61K007/09 A61K007/06.

---

☐ 121. 5229104. 29 Apr 91; 20 Jul 93. Artificial tanning compositions containing positively charged paucilamellar vesicles. Sottery; John P., et al. 424/59; 424/195.18 424/450 424/60 424/63 424/757 514/167 514/251 514/458 514/474 514/725 514/847 514/938. A61K007/42 A61K007/44 A61K007/48 A61K009/10.

---

☐ 122. 5189066. 19 Jul 91; 23 Feb 93. Pharmaceutical compositions of tebufelone. Kelm; Gary R., et al. 514/678; 514/689 514/941 514/960 514/961. A61K031/12.

---

☐ 123. 4981681. 26 Aug 88; 01 Jan 91. Lotion mixture and method of treating psoriasis. Tosti; Vittorio. 514/161; 424/78.05 514/159 514/337 514/356 514/458 514/560 514/848 514/863. A61K031/685 A61K007/48.

---

☐ 124. JP403176420A. 22 Nov 90. 31 Jul 91. MEDICINAL COMPOSITION OF TEBUFELONE. KELM, GARY R, et al. A61K031/12; A61K009/08 A61K009/48 A61K031/12 A61K031/12 A61K047/34.

---

☐ 125. JP361225121A. 29 Mar 85. 06 Oct 86. PREVENTIVE FOR ASTHMA AND PREPARATION THEREOF. ABE, TORU, et al. A61K031/41; A61K009/00 A61K009/14 A61K009/16 A61K009/20 A61K009/48 A61K009/70 A61K047/00 C07D257/04.

---

☐ 126. WO000166109A1. 07 Mar 01. 13 Sep 01. ORAL PHARMACEUTICAL COMPOSITIONS CONTAINING N-SULPHONYLINDOLINE DERIVATIVES. BLUNDELL, ROSS, et al. A61K031/40; A61K009/48.

---

☐ 127. DE019946184A1. 21 Sep 99. 29 Mar 01. Retinoid-containing emulsions with continuous water phase, for topical application e.g. for skin treatment, repair or renewal, having structured, polar, viscous oil phase to stabilize active agent. JENNING, VOLKHARD. A61K031/07; A61P017/00.

---

☐ 128. FR002776189A1. 09 Jun 99. 24 Sep 99. Oral compositions having improved absorption in the gastro-intestinal tract containing a polyglycolized glyceride. SASLAWSKI, OLIVIER, et al. A61K009/20; A61K031/155.

---

☐ 129. RU 2242216C. Lifting-cream for dry and sensitive skin. ALAVERDIEV, I M O, et al. A61K007/00 A61K007/48.

---

☐ 130. KR2004088794A. Eye drops composition containing trehalose and fluran used as artificial tear solution, therapeutic for eye dryness, contact lens solution and drug delivery material. KIM, B C.

A61K031/719.

---

☐ 131. US20040087460A. Dishwashing cleaning wipe for dishware, flatware, pots and pans, contains water insoluble substrate impregnated with liquid cleaning composition comprising at least one anionic surfactant and water. KELLY, A R, et al. A47L013/17 B32B023/10 C11D001/83 C11D001/86 C11D003/20 C11D003/34 C11D017/00 C11D017/04.

---

☐ 132. US 6727210B. Cleansing composition for use in cleaning epidermis of user comprises e.g. water, aloe extract, citric acid, disodium ethylenediaminetetraacetic acid, potassium sorbate, propylene glycol, quaternium-52 and benzethonium chloride. PERDEW, D E. C11D017/00.

---

☐ 133. US20030203831A. Dishwashing cleaning wipe for dishwasher, fabric, hair, teeth or human skin, comprises liquid cleaning composition coated and impregnated into water insoluble substrate, and comprising anionic surfactant, surfactant and water. THOMAS, B, et al. C11D017/00.

---

☐ 134. US 6534472B. Dishwashing cleaning wipe for, e.g. dishware, flatware, pots and pans, comprises water insoluble substrate impregnated with antibacterial cleaning composition. ARVANITIDOU, E, et al. C11D001/94 C11D003/02 C11D003/20 C11D003/37 C11D017/00 C11D017/04.

---

☐ 135. WO2002100418A. Fluid composition for treating ectoparasite infestation, comprising particulate water insoluble inorganic substance which reacts exothermically with water to create localized temperatures. SCHLESINGER, M S, et al. A01K013/00 A01M001/00 A01M007/00 A01N025/00 A01N025/20 A01N059/00 A01N059/06 A61F007/00 A61K000/00 A61K007/00 A61K007/06 A61K033/00 A61K033/06 A61N001/00 A61P031/12 A61P031/14 A61P033/14.

---

☐ 136. US 6489284B. Dishwashing cleaning wipe comprises water-insoluble substrate impregnated with cleaning composition comprising anionic surfactant, alkyl polyglucoside surfactant, alkyl monoalkanol amide, alkanol and water. BARRY, D, et al. A47L013/17 C11D001/83 C11D001/86 C11D003/20 C11D003/34 C11D017/00 C11D017/04.

---

☐ 137. US20020164381A. Treatment of infestations of cutaneous mites e.g. harvest mites involves applying a composition comprising sulfur, its derivative and a carrier. GANS, E H, et al. A01N025/32 A61K000/00 A61K007/75 A61K033/04.

---

☐ 138. WO 200209637A. Composition useful for the treatment of e.g. mucositis and stomatitis comprises hyaluronic acid, glycyrrhetic acid and polyvinylpyrrolidone in mixture with excipients and adjuvants. MASTRODONATO, M, et al. A61K000/00 A61K009/14 A61K031/00 A61K031/19 A61K031/715 A61K031/728 A61K031/79 A61K047/10 A61K047/12 A61K047/18 A61K047/22 A61K047/32 A61K047/36 A61K047/38 A61K047/42 A61P001/02 A61P001/04 A61P015/02 A61P019/08 A61P029/00 A61K031/79 A61K031:19 A61K031:728 A61K031/79 A61K031:19 A61K031:728 A61K031/79 A61K031:19 A61K031:728.

---

☐ 139. US20010048931A. Oral megestrol acetate formulation comprises a stable flocculated aqueous suspension containing a polyol and surfactant. CHAO, J C, et al. A61K009/00 A61K031/56.

---

☐ 140. CN 1255568A. Surface modifier of polypropylene fibre for cigarettes and its surface modifying process. CHEN, Z, et al. A24D003/02 D06M015/263.

---

☐ 141. US 5945098A. A storage stable, intravenous-administrable immunoglobulin composition, useful in immunotherapy and in diagnostic processes. GRAF, C R, et al. A61K038/21.

---

- ☐ 142. WO 9703655A. Soft gelatin capsules contg. compsn. comprising NSAID - which is dissolved in di:methyl- isosorbide or a mixt. of this with, e.g., poly:sorbate, polyethylene glycol or food stock oils. MUNDEN, J W. A61K009/48 A61K031/19 A61K031/192 A61K031/40 A61K031/54 A61K047/10 A61K047/14 A61K047/26 A61K047/34 A61K047/36.
- ☐ 143. WO 9640217A. Stable compsn. increasing proliferation of cells of myeloid lineage - contains thrombopoietin, surface adsorption inhibitor and physiologically acceptable buffer and salt. KONTOR, K M, et al. A61K009/08 A61K037/02 A61K038/00 A61K038/19 A61K038/22 A61K047/16 A61P007/06 A61P043/00 C07K014/52.
- ☐ 144. DE 4327327A. Nonionic surfactant compsns. - comprising n-alkyl polyoxyethylene-polyoxyp-ropylene ether and other nonionic surfactant. FABRY, B, et al. B01F017/00 B03D001/00 C11D001/722 C11D001/825 C23G001/24.
- ☐ 145. EP 605203A. Antiallergic compsn. for ophthalmic or nasal use contg. cetirizine - also opt. contains cyclodextrin cpd. surfactant and/or water soluble polymer, for treating e.g. allergic or vernal conjunctivitis. IKEJIRI, Y, et al. A61K009/08 A61K009/12 A61K031/445 A61K031/495 A61K047/40 A61K047/48 A61P011/02 A61P027/02 C07D295/08.
- ☐ 146. US 5304370A. Human hair relaxer compsn. - of mixed emulsions contg sodium hydroxide surfactant and oil. HAWKINS, G R, et al. A61K007/06 A61K007/09.
- ☐ 147. DE 4229442A. Dispersant contg. alkyl:poly:glycoside or alkenyl:poly:glycoside - and poly-hydroxy fatty acid amide, esp. for use in prepn. of pigment and dye compsns.. FABRY, B, et al. A01N003/00 B01F017/00 B01F017/22 B01F017/56 C07C233/18 C07H015/04 C09B067/20 C09B067/42 C09B067/46 D06P001/48 D06P001/649.
- ☐ 148. WO 9323515A. Colour stabilisation of surfactant with amino-alkanoic acid cpds. - which chelate iron and are readily biodegradable, useful with anionic, nonionic and amphoteric surfactant. BLUM, H, et al. A61K007/00 A61K007/075 A61K007/50 B01F017/00 C07C229/04 C09K015/20 C11D003/20 C11D003/30 C11D003/33.
- ☐ 149. RD 353036A. Stabilisation of liq. enzyme preparations for contact lens rinse - by addn. of boric acid or borate ester to improve enzyme stability. A61L000/00.
- ☐ 150. EP 522936A. Compsn. for treating cancer, tumour and leukaemia - comprises taxan in soln. in surfactant with minimal ethanol. BASTART, J, et al. A61K000/00 A61K009/00 A61K009/08 A61K009/50 A61K031/33 A61K031/335 A61K031/337 A61K031/34 A61K047/010 A61K047/026 A61K047/044 A61K047/10 A61K047/14 A61K047/26 A61K047/44 C07D305/14.

[Generate Collection](#)
[Print](#)

Terms	Documents
L5 same (supposit\$ or composition or vaccine or insertion or application or applying).ti,ab,clm.	160

[Prev Page](#)
[Next Page](#)



US005945098A

**United States Patent** [19]

Sarno et al.

[11] Patent Number: **5,945,098**[45] Date of Patent: **\*Aug. 31, 1999**[54] **STABLE INTRAVENOUSLY-ADMINISTRABLE IMMUNE GLOBULIN PREPARATION**[75] Inventors: **Maria Erlinda C. Sarno, Cerritos; Rodolfo Anthony Vasquez, Norwalk; Sau-Gee Yung, Rialto; Clifford R. Graf, Lakeview Terrace, all of Calif.**[73] Assignee: **Baxter International Inc., Deerfield, Ill.**

[\*] Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

[21] Appl. No.: **08/935,294**[22] Filed: **Sep. 22, 1997****Related U.S. Application Data**

[63] Continuation of application No. 08/504,854, Jul. 20, 1995, abandoned, which is a continuation of application No. 08/317,214, Oct. 3, 1994, abandoned, which is a continuation of application No. 08/178,432, Jan. 6, 1994, abandoned, which is a continuation of application No. 07/866,089, Apr. 6, 1992, abandoned, which is a continuation of application No. 07/473,554, Feb. 1, 1990, abandoned.

[51] Int. Cl.<sup>6</sup> ..... **A61K 38/21**[52] U.S. Cl. .... **424/85.5; 514/2; 514/12; 514/21; 530/387.1; 530/389.1; 530/380; 530/390.5**[58] Field of Search ..... **424/85.5; 514/2, 514/12, 21; 530/387.1, 389.1, 380, 390.5**[56] **References Cited****U.S. PATENT DOCUMENTS**

3,903,262	9/1975	Pappenhagen et al.	424/85
4,165,370	8/1979	Coval	424/85.8
4,168,303	9/1979	Nishida et al.	530/387
4,186,192	1/1980	Lundblad et al.	424/85.8
4,362,661	12/1982	Ono et al.	424/85.1
4,371,520	2/1983	Uemura et al.	424/85.8
4,374,763	2/1983	Takagi	424/85.8
4,384,993	5/1983	Sato et al.	424/85.1
4,396,608	8/1983	Tenold	424/85.8
4,439,421	3/1984	Hooper et al.	424/85
4,477,432	10/1984	Hardie	424/85.1
4,482,483	11/1984	Curry et al.	530/387
4,540,573	9/1985	Neurath et al.	424/85
4,565,651	1/1986	Ohmura et al.	514/8
4,597,966	7/1986	Zolton et al.	424/85.8
4,613,501	9/1986	Horowitz	424/89
4,623,717	11/1986	Fernandes et al.	530/381
4,637,834	1/1987	Thurrow	514/3
4,639,513	1/1987	Hou et al.	530/387

4,692,331	9/1987	Uemura et al.	424/85
4,719,290	1/1988	Curry et al.	530/357
4,721,777	1/1988	Uemura et al.	530/388.15
4,764,369	8/1988	Neurath et al.	424/89
4,789,545	12/1988	Woods et al.	424/101
4,876,088	10/1989	Hirao et al.	424/85.8
4,877,608	10/1989	Lee et al.	424/85.8
5,177,194	1/1993	Saras et al.	530/412

**FOREIGN PATENT DOCUMENTS**

0187712	7/1986	European Pat. Off.
0196761	10/1986	European Pat. Off.
2500076	7/1976	Germany
53-47515	4/1978	Japan
55-164630	12/1980	Japan
59-97057	6/1984	Japan

**OTHER PUBLICATIONS**

Tomono et al., *Vox Sang* vol. 51 (2) pp. 81-86 (1986).

Stephan, W., "Undegraded Gamma-Globulin for Intravenous Therapy: A New Preparation of Immune Serum Globulin for Intravenous Administration," *XXIVth Scientific Meeting of the Blood Research Institute*, pp. 469-478, 1971.

Fernandes et al., "Preparation of a Stable Intravenous Gamma-Globulin: Process Design and Scale-Up," *Vox Sang*. 39: 101-12 (1980).

Gerber, "Stabilization of Gamma Globulin with Copper Complexes: Possible Relevance to the Etiology of Rheumatoid Arthritis," *Inflammation Diseases and Copper* (1982).

Gerber, "Inhibition of Denaturation of Human Gamma Globulin by a Mixture of L-Histidine, L-Cystine, and Copper, and Its Clinical Implication in Rheumatoid Arthritis," *Arthritis and Rheumatism*, vol. 19, 3:593-601 (1976).

Florence et al., "Emulsion Stabilization by Non-Ionic Surfactants: Experiment and Theory," *J. Pharm. Pharmac.* 23:153-169 (1971).

Miller, G., et al., "Stabilization of Cytotoxic Antibody and Complement with Cysteine," *J. of Immun.*, vol. 101, No. 5, pp. 1074-1077 (1968).

**Primary Examiner**—Chhaya D. Sayala**Attorney, Agent, or Firm**—Rothwell, Figg, Ernst & Kurz, p.c.[57] **ABSTRACT**

Stable, intravenously-administrable immune globulin preparations are stabilized against aggregation and polymerization and rendered isotonic with amino acid(s) and non-ionic detergents, polysorbate and polyethylene glycol. The immunoglobulins are derived from human or animal sources, or from hybridomas. Optional, additional stabilizers include various physiologically-acceptable carbohydrates and salts. Polyvinylpyrrolidone can be used in addition to the amino acid(s). Apart from the immunoglobulin itself, the preparations are otherwise essentially protein free. The preparations are useful in immunotherapy and as diagnostic reagents.

**15 Claims, No Drawings**

US-PAT-NO: 5643587

DOCUMENT-IDENTIFIER: US 5643587 A

TITLE: Composition and method for under-eye skin lightening

DATE-ISSUED: July 1, 1997

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Scancarella; Neil D.	Wyckoff	NJ		
Duffy; John A.	West Milford	NJ		
Garrison; Mark S.	White Plains	NY		
Menon; Gopinathan K.	Wayne	NJ		

US-CL-CURRENT: 424/401; 424/195.16, 424/62

## CLAIMS:

What is claimed is:

1. A composition for topical treatment of skin discolorations found below the eyes, comprising:

- (a) about 0.5 to about 10.0% live yeast cell extract,
- (b) about 0.5 to about 10.0% magnesium ascorbyl phosphate,
- (c) about 0.1 to about 5.0% tocopherol acetate,
- (d) about 0.01 to about 1.0% retinol palmitate, and
- (e) a vehicle which maintains active ingredient levels.

2. The composition of claim 1, comprising:

- (a) about 8.5% emollient oils,
- (b) about 7.5% waxes,
- (c) about 5.0% glycerin,
- (d) about 3.52% live yeast cell extract,
- (e) about 3.0% magnesium ascorbyl phosphate,
- (f) about 1.0 tocopherol acetate,
- (g) about 0.5% retinol palmitate,
- (h) about 0.7% vitamins,



- (i) about 2.5% emulsifiers, and
- (j) about 0.35% thickeners.

3. The composition of claim 2, wherein said emollient oils are selected from the group consisting of dioctyl maleate, grape and sunflower seed oils, and squalane; said waxes are selected from the group consisting of soya sterols, glyceryl monostearate, cetyl alcohol and myristyl myristate; said vitamins is vitamin K; said emulsifiers are selected from the group consisting of PEG 40 stearate, PEG 24 cholsteryl ether and polysorbate 20; and said thickeners are selected from the group consisting of xantham gum, Carbopols 934 and 941 and further comprising ascorbyl palmitate.

4. The composition of claim 1, comprising:

- (a) about 10.0% emollient oils,
- (b) about 5.0% glycerin,
- (c) about 4.75% waxes,
- (d) about 3.5% magnesium ascorbyl phosphate,
- (e) about 3.25% live yeast cell extract,
- (f) about 1.0 tocopherol acetate,
- (g) about 0.25% retinol palmitate,
- (h) about 0.8% vitamins,
- (i) about 1.75% emulsifiers, and
- (j) about 0.35% thickeners.

5. The composition of claim 4, wherein said emollient oils are selected from the group consisting of dioctyl maleate, grape and sunflower seed oils, and squalane; said waxes are selected from the group consisting of soya sterols, glyceryl monostearate, cetyl alcohol and myristyl myristate; said vitamins is vitamin K; said emulsifiers are selected from the group consisting of PEG 40 stearate, PEG 24 cholsteryl ether and polysorbate 20; and said thickeners are selected from the group consisting of xantham gum, Carbopols 934 and 941 and further comprising ascorbyl palmitate.

6. The composition of claim 1, comprising:

- (a) about 1.5% emollient oils,
- (b) about 1.5% glycerin,
- (c) about 5.0% live yeast cell extract,
- (d) about 2.0% magnesium ascorbyl phosphate,

- (e) about 0.1 tocopherol acetate,
- (f) about 0.1% retinol palmitate,
- (g) about 0.2% vitamins,
- (h) about 1.5% emulsifier, and
- (j) about 0.75% thickeners.

7. The composition of claim 6, wherein said emollient oils are selected from the group consisting of dioctyl maleate, grape and sunflower seed oils, and squalane; said vitamins is vitamin K; said emulsifiers are selected from the group consisting of PEG 40 stearate, PEG 24 cholsteryl ether and polysorbate 20; and said thickeners are selected from the group consisting of xantham gum, Carbopols 934 and 941 and further comprising ascorbyl palmitate.

8. A method for topical treatment of skin discolorations found below the eyes comprising the application of a composition comprised of:

- (a) about 0.5 to about 10.0% live yeast cell extract,
- (b) about 0.5 to about 10.0% magnesium ascorbyl phosphate,
- (c) about 0.09 to about 0.6% retinol palmitate,
- (d) about 0.1 to about 5.0% tocopherol acetate, and
- (e) a vehicle which maintains active ingredient levels.

9. The method of claim 8, wherein said composition comprises:

- (a) about 8.5% emollient oils,
- (b) about 7.5% waxes,
- (c) about 5.0% glycerin,
- (d) about 3.52% live yeast cell extract,
- (e) about 3.0% magnesium ascorbyl phosphate,
- (f) about 1.0 tocopherol acetate,
- (g) about 0.5% retinol palmitate,
- (h) about 0.7% vitamins,
- (i) about 2.5% emulsifiers, and
- (j) about 0.35% thickeners.

10. The method of claim 9, wherein said emollient oils are selected from the

group consisting of dioctyl maleate, grape and sunflower seed oils, and squalane; said waxes are selected from the group consisting of soya sterols, glyceryl monostearate, cetyl alcohol and myristyl myristate; said vitamins is vitamin K; said emulsifiers are selected from the group consisting of PEG 40 stearate, PEG 24 cholsteryl ether and polysorbate 20; and said thickeners are selected from the group consisting of xantham gum, Carbopols 934 and 941 and further comprising ascorbyl palmitate.

11. The method of claim 8, wherein said composition comprises:

- (a) about 10.0% emollient oils,
- (b) about 5.0% glycerin,
- (c) about 4.75% waxes,
- (d) about 3.5% magnesium ascorbyl phosphate,
- (e) about 3.25% live yeast cell extract,
- (f) about 1.0 tocopherol acetate,
- (g) about 0.25% retinol palmitate,
- (h) about 0.8% vitamins,
- (i) about 1.75% emulsifiers, and
- (j) about 0.35% thickeners.

12. The method of claim 11, wherein said emollient oils are selected from the group consisting of dioctyl maleate, grape and sunflower seed oils, and squalane; said waxes are selected from the group consisting of soya sterols, glyceryl monostearate, cetyl alcohol and myristyl myristate; said vitamins is vitamin K; said emulsifiers are selected from the group consisting of PEG 40 stearate, PEG 24 cholsteryl ether and polysorbate 20; and said thickeners are selected from the group consisting of xantham gum, Carbopols 934 and 941 and further comprising ascorbyl palmitate.

13. The method of claim 8, wherein said composition comprises:

- (a) about 1.5% emollient oils,
- (b) about 1.5% glycerin,
- (c) about 5.0% live yeast cell extract,
- (d) about 2.0% magnesium ascorbyl phosphate,
- (e) about 0.1 tocopherol acetate,
- (f) about 0.1% retinol palmitate,
- (g) about 0.2% vitamins and vitamin derivatives,

(h) about 1.5% emulsifier, and

(j) about 0.75% thickeners.

14. The method of claim 13, wherein said emollient oils are selected from the group consisting of dioctyl maleate, grape and sunflower seed oils, and squalane; said vitamins is vitamin K; said emulsifiers are selected from the group consisting of PEG 40 stearate, PEG 24 cholsteryl ether and polysorbate 20; and said thickeners are selected from the group consisting of xantham gum, Carbopols 934 and 941 and further comprising ascorbyl palmitate.

15. A composition for topical treatment of skin discolorations found below the eyes, comprising:

(a) about 0.5 to about 10.0% live yeast cell extract,

(b) about 0.5 to about 10.0% magnesium ascorbyl phosphate, and

(c) a vehicle which maintains active ingredient levels.

16. The composition of claim 15, further comprising about 0.1 to about 5.0% tocopherol acetate.

17. The composition of claim 15, further comprising about 0.01 to about 1.0% retinol palmitate.

18. A method for topical treatment

First Hit

L8: Entry 141 of 160

File: DWPI

Aug 31, 1999

DERWENT-ACC-NO: 1999-539002

DERWENT-WEEK: 199945

COPYRIGHT 2005 DERWENT INFORMATION LTD

TITLE: A storage stable, intravenous administrable immunoglobulin composition, useful in immunotherapy and in diagnostic processes

Basic Abstract Text (2):

DETAILED DESCRIPTION - A storage stable, intravenously-administrable immune globulin composition (I) comprises an aqueous solution of immune globulin, 0.1-0.3 M glycine, 0.0005-0.01% (w/v) polysorbate, and less than 0.2 g polyethylene glycol (PEG). (I) is essentially protein-free apart from the immune globulin.

First Hit

L8: Entry 158 of 160

File: DWPI

Oct 29, 1986

DERWENT-ACC-NO: 1986-286406

DERWENT-WEEK: 199718

COPYRIGHT 2005 DERWENT INFORMATION LTD

TITLE: Recovering pure protein from cultures of genetically modified yeast - by lysis in presence of nonionic detergent then pptn. of contaminants with polyethylene glycol

INVENTOR: GILLES, D; SIMONET, G ; VANWIJNEN, F

PATENT-ASSIGNEE: SMITH KLINE-RIT (SMIT), SMITH KLINE-RIT SA (SMIK)

PRIORITY-DATA: 1987US-0042914 (April 27, 1987), 1985US-0719601 (April 3, 1985)

Search Selected

Search All

Clear

## PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
<input type="checkbox"/> EP 199698 A	October 29, 1986	F	015	
<input type="checkbox"/> PT 82219 A	September 16, 1986		000	
<input type="checkbox"/> AU 8653786 A	October 9, 1986		000	
<input type="checkbox"/> NO 8601234 A	October 27, 1986		000	
<input type="checkbox"/> JP 61249398 A	November 6, 1986		000	
<input type="checkbox"/> FI 8601251 A	October 4, 1986		000	
<input type="checkbox"/> DK 8601522 A	October 4, 1986		000	
<input type="checkbox"/> HU 40810 T	February 27, 1987		000	
<input type="checkbox"/> ZA 8602374 A	May 18, 1987		000	
<input type="checkbox"/> CN 8601798 A	October 15, 1986		000	
<input type="checkbox"/> US 4683294 A	July 28, 1987		000	
<input type="checkbox"/> ES 8706166 A	August 16, 1987		000	
<input type="checkbox"/> DD 247685 A	July 15, 1987		000	
<input type="checkbox"/> CS 8602233 A	February 15, 1988		000	
<input type="checkbox"/> RO 95349 A	September 15, 1988		000	
<input type="checkbox"/> US 4857317 A	August 15, 1989		000	
<input type="checkbox"/> CA 1272457 A	August 7, 1990		000	
<input type="checkbox"/> IL 78824 A	April 15, 1991		000	
<input type="checkbox"/> SU 1604144 A	October 30, 1990		000	
<input type="checkbox"/> EP 199698 B	August 28, 1991		000	
<input type="checkbox"/> DE 3681058 G	October 2, 1991		000	

<input type="checkbox"/> JP 2512430 B2	July 3, 1996	008	C12P021/00
<input type="checkbox"/> KR 9410283 B1	October 22, 1994	000	C12N015/00

DESIGNATED-STATES: AT BE CH DE FR GB IT LI LU NL SE AT BE CH DE FR GB IT LI LU NL SE

CITED-DOCUMENTS: 4.Jnl.Ref; A3...198921 ; EP 106828 ; No-SR.Pub ; US 4144130 ; 5.Jnl.Ref

## APPLICATION-DATA:

PUB-NO	APPL-DATE	APPL-NO	DESCRIPTOR
EP 199698A	April 2, 1986	1986EP-0870041	
JP 61249398A	April 2, 1986	1986JP-0097518	
ZA 8602374A	April 1, 1986	1986ZA-0002374	
US 4683294A	April 3, 1985	1985US-0719601	
ES 8706166A	April 2, 1986	1986ES-0553633	
US 4857317A	April 27, 1987	1987US-0042914	
SU 1604144A	April 2, 1986	1986SU-4027296	
JP 2512430B2	April 2, 1986	1986JP-0077518	
JP 2512430B2		JP 61249398	Previous Publ.
KR 9410283B1	April 2, 1986	1986KR-0002480	

INT-CL (IPC): A23J 1/18; A61K 37/02; A61K 39/29; C07C 3/12; C07K 1/14; C07K 3/00; C07K 14/02; C07K 15/04; C12N 15/00; C12P 21/00; C12P 21/02; C12P 21/00; C12R 1/865

ABSTRACTED-PUB-NO: EP 199698A

## BASIC-ABSTRACT:

A protein (I) bound to cells is extd. and purified from the supernatant of a genetically-modified yeast cell culture which has been lysed in presence of a nonionic detergent, comprises first adjusting the supernatant to pH 5.9-6.1. Then liq. or solid polyethylene glycol (PEG) is added until the supernatant becomes clear, then this treated with (1) a bivalent metal cation or (2), opt. after ultrafiltration, with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> to separate (I). Pref. liq. PEG is added to final concn. 10-35 vol.% or solid PEG to 6-12wt.%.

Pref. the clarified supernatant is treated with Ca or Mn ions to a final concn. of 80 mm. The ppte. is removed and the supernatant purified by ultrafiltration, gel permeation of the retentate and ion-exchange chromatography of the (I)-contg. peak. Opt. (I) is further purified by a second gel permeation step and centrifuging in a CsCl gradient. Alternatively, the clarified supernatant is mixed with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> to 40-50% satn. and (I) recovered in the PEG phase. This phase can be ultrafiltered and then subjected to gel permeation and ion-exchange chromatography. The nonionic detergent is esp. polysorbate 20.

USE/ADVANTAGE - The method is esp. applied to hepatitis B surface antigen (recovered in the form of micelles with a polysorbate, useful for making vaccines) or alpha-1-antitrypsin. Treatment with PEG provides efficient pptn. of most of the contaminants (typically 75% of proteins; 90% of polysaccharides, 94% of nucleic acids and 45% of lipids), while (I) remains in soln. A relatively high yield of pure (I) is achieved.

ABSTRACTED-PUB-NO: EP 199698B  
EQUIVALENT-ABSTRACTS:

A process for the extraction and purification of a protein, bound to the cell, from the supernatant of a culture of genetically manipulated yeast cells which have produced said protein and burst in the presence of a non-ionic detergent, characterised in that the pH of the supernatant is adjusted to 6 (+/-0.1), liquid or solid polyethylene glycol is added until said supernatant has been clarified, and the clarified supernatant is treated either with a divalent metal cation, or after ultrafiltration if appropriate, with ammonium sulphate in order to separate out said protein. (19pp)

US 4683294A

Improved extn. and purificn. of cell-bound protein obtd. from the supernatant of engineered yeast cells producing such protein, comprises disruption using a nonionic detergent.

Process comprises (a) adjusting the supernatant to pH 6(+/-0.1); (b) adding liq. or solid polyethylene glycol up to the clarification of the supernatant; and (c) treating clarified prod. with a bivalent metal cation or with ammonium sulphate for sepg. the protein before or after ultrafiltration.

Pref. final concn. of liq. polyethylene glycol is 10-35% (v/v) and corresp. solid is 6-12% (w/v).

USE - For extn. and purificn. of hepatitis B surface antigen or alpha-1-antitrypsin. (7pp)d

US 4857317A

Hepatitis B surface antigen/polysorbate composite micelle contains 15-35% (w/w) polysorbate, as active ingredient in corresp. vaccine. Prodn. comprises (a) prepg. a supernatant of engineered yeast cells disrupted using a non-ionic detergent; (b) pptg. cqntominants by polyethylene glycol; and (c) treating supernatant obtd. with a bivalent metal cation, or with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> after eventual ultrafiltration. Hepatitis B virus surface antigen and alpha-1-antitrypsin are each thus extracted and purified.

USE - To protecting human against hepatitis B virus infection by administering intramuscularly. (7pp)

CHOSEN-DRAWING: Dwg.0/0

DERWENT-CLASS: A96 B04 D16

CPI-CODES: A05-H02; A12-W11L; B02-V02; B04-B04A5; B04-B04C1; D05-H07;





Europäisches Patentamt  
European Patent Office  
Office européen des brevets

Numéro de publication:

0 199 698  
A2



# DEMANDE DE BREVET EUROPEEN

Numéro de dépôt: 86870041.0

Int. Cl.<sup>4</sup> C12P 21/02, C12N15 00

Date de dépôt: 02.04.86

30 Priorité: 03.04.85 US 719601

43 Date de publication de la demande:  
29.10.86 Bulletin 86/44

64 Etats contractants désignés:  
AT BE CH DE FR GB IT LI LU NL SE

71 Demandeur: Smith Kline - RIT Société  
anonyme dite:  
rue du Tilleul, 13  
B-1320 Genval (Rixensart)(BE)

72 Inventeur: van Wijnendaele, Frans  
Teckerstraat, 12  
B-3052 Ottenburg(BE)  
Inventeur: Gilles, Daniel  
Rue Mascau, 13  
B-1320 Genval(BE)  
Inventeur: Simonet, Guy  
Chaussée de Wavre, 14  
B-5920 Perwez(BE)

74 Mandataire: Tasset, Gérard  
SMITHKLINE - RIT rue de l'Institut, 89  
B-1330 Rixensart(BE)

54 Procédé d'extraction et de purification de protéines issues de milieux de culture les produisant.

57 Le procédé s'applique au surnageant de cellules de levure produites génétiquement et éclatées en présence d'un détergent non ionique; il comprend la précipitation des contaminants par le polyéthylène glycol et le traitement de ce surnageant par un cation métallique bivalent ou, après ultrafiltration éventuelle, par le sulfate d'ammonium.

EP 0 199 698 A2

# Procédé d'extraction et de purification de protéines issues de milieux de culture les produisant

## Domaine de l'invention

La présente invention a pour objet un procédé perfectionné d'extraction et de purification de protéines issues de milieux de culture les produisant; plus particulièrement, le procédé concerne l'extraction et la purification de l'antigène de surface du virus de l'hépatite B et de l'alpha-1-antitrypsine produits par des techniques d'ADN recombinant dans des cultures cellulaires produites génétiquement, plus particulièrement des cultures de souches de levure produites génétiquement.

## Fondement de l'invention

Divers microorganismes, parmi lesquels les bactéries et les levures, peuvent être utilisés comme organismes hôtes pour différents plasmides contenant une molécule d'ADN renfermant une séquence nucléotidique codant pour une protéine spécifique. Parmi ces microorganismes, les levures sont actuellement préférées et d'un emploi courant. Par exemple, la demande de brevet européen publiée sous le numéro 0 106 828 décrit l'utilisation des souches de *Saccharomyces cerevisiae* pour la production de l'antigène de surface du virus de l'hépatite B (HBsAg) et la demande de brevet européen publiée sous le numéro 0 103 409 est relative à la production de l'alpha-1-antitrypsine au départ de plasmides de levure.

La production de protéines dans des souches de levure présente, en effet, de nets avantages par rapport à la production dans des souches bactériennes. Ces avantages proviennent de la croissance plus aisée des levures dans des fermenteurs à grande échelle et du fait que, contrairement aux bactéries, les levures ressemblent assez bien aux cellules de mammifères pour leur capacité d'additionner des groupes glucidiques aux protéines qui viennent d'être synthétisées.

Toutefois, l'extraction et la purification de la protéine issue d'une culture de levure donnent lieu à des problèmes techniques dus à la composition chimique plutôt complexe de la cellule de levure et plus particulièrement à cause de la présence des taux lipidiques élevés lorsqu'on prolonge la croissance de la levure pour améliorer le rendement de la production en microcystides.

Par exemple, à cet égard, le *Saccharomyces* est supposée se composer de 3 couches: (a) une couche interne de  $\beta$ -glucane insoluble en milieu alcalin, (b) une couche centrale de  $\beta$ -glucane alcalino-soluble et (c) une couche externe de glycoprotéine dans laquelle le glucide se compose de

mannane phosphorylé; en dessous de la paroi se trouve une membrane cytoplasmique qui se compose d'un mélange très complexe constitué de lipides neutres (mono-, di- et triglycérides), de stéroïdes libres et estérifiés, d'un soningolipide complexe, de glycérophosphatides et de glycolipides neutres et acides; le noyau renferme de l'ADN, diverses espèces d'ARN et un polyphosphate; les vacuoles peuvent contenir une grande variété de composés de poids moléculaires élevés ou faibles; elles servent de vésicules de réserve destinées à de nombreuses hydrolases; les mitochondries sont riches en composés lipidiques, phospholipidiques et ergostéroliques en provenance du système membraneux et le cytoplasme contient entre autres de grandes quantités de ribosomes, de polyphosphates, de glycogènes et de nombreux enzymes de la glycolyse. La plupart des cellules de levure (telle que l'espèce *Saccharomyces*) contiennent également une certaine quantité de lipide sous la forme de globules dont la quantité augmente dans les cultures prolongées.

De nombreux procédés à étapes multiples ont été décrits pour l'extraction et la purification de protéines issues de différentes sources. Des exemples de production de protéines par des microorganismes obtenus génétiquement sont ceux publiés par Th. STAHELIN et al. (J. Biol. Chem. 256:9750-54; 1981), K. MURRAY et al. (The Embo J. 3:645-650; 1984) et R.A. HITZEMAN et al. (Nucl. Ac. Res. 11:2745-2763; 1983).

En fait, lorsque la protéine produite est liée à la cellule, ces différents procédés impliquent 3 séries d'étapes.

Dans la première série d'étapes, la protéine désirée est isolée de l'intérieur de la cellule. A cette fin, les cellules sont soit lysées (p.ex. par traitement enzymatique) ou éclatées (p.ex. par des forces mécaniques telles que les forces de cisaillement (p.ex. la cellule de pression X ou la cellule de pression de French) ou agitées avec des billes de verre, avec addition éventuelle d'un détergent (voir par exemple K. MURRAY et al., loc. cit. et R.A. HITZEMAN et al., loc. cit.).

Dans la deuxième série d'étapes, le milieu est enrichi en protéine désirée, p.ex. par précipitation fractionnée par ajout de sulfate d'ammonium et/ou en présence de polyéthylène glycol.

Enfin, dans la troisième série d'étapes, tous les contaminants sont effectivement éliminés du milieu, p.ex. par une ou plusieurs opérations du genre ultrafiltration, chromatographie par échange d'ions et par filtration sur gel et centrifugation en gradient isopycnique.

Dans un tel procédé, il est évident que les contaminants accompagnant les protéines (cités par R.A. HITZEMAN et al., loc. cit.), les acides nucléiques et les lipides et plus particulièrement les taux de lipides élevés exercent une forte influence sur au moins une des étapes de la troisième série (p.e. l'ultrafiltration et la chromatographie sur colonne) et que les hydrolases doivent être rapidement éliminées du milieu. De plus, il a également été observé que la précipitation au sulfate d'ammonium n'est pas possible sans une délipidation préalable, étant donné que les lipides interfèrent dans cette précipitation.

En conséquence, il est de la première importance de pouvoir disposer d'une méthode par laquelle on puisse éliminer la plupart des contaminants dès le début de la troisième série d'étapes.

Parmi les procédés déjà décrits, certains citent l'utilisation du polyéthylène glycol comme agent de précipitation sélectif des protéines et un procédé de précipitation des lipoprotéines à partir du plasma en faisant appel aux interactions lipoprotéine-polyanionmétal a également été signalé.

La méthode de précipitation fractionnée des protéines à l'aide de polymères non ioniques hydrosolubles, en particulier le polyéthylène glycol (PEG) a été introduite par POLSON et al. (Biochem. Biophys. Acta 82:463-475; 1964) et discutée par divers auteurs. Parmi eux, W. HONIG et al. (Analyt. Biochem. 72:502-512; 1976) mentionnent que "la spécificité de la précipitation, c'est-à-dire le rapport de la protéine désirée et de la protéine totale peut être amélioré en utilisant des fractions de PEG d'un poids moléculaire moyen inférieur au PEG 6000 généralement employé". Néanmoins, quoique "des concentrés de protéines plasmatiques individuelles puissent être obtenus en agissant sur le pH", les auteurs ajoutent cependant que "la purification de mélanges protéiniques plus complexes tels que le surnageant d'un homogénat de cellule est considérablement plus médiocre".

P.R. FOSTER et al. (Biochim. Biophys. Acta 317: 505-516; 1973) ont décrit une méthode de précipitation d'enzymes à partir d'extraits cellulaires de *Saccharomyces cerevisiae* au moyen du PEG. Des méthodes de concentration et de purification de virus et de bactériophages à l'aide de PEG ont été publiées par B.P. VAJDA (Folia Microbiol. 23: 88-96; 1978) et G.J. LANCZ (Arch. Virology, 42:303-306; 1973). Dans le domaine de l'isolement de l'antigène de l'hépatite, la purification de l'antigène de surface de l'hépatite B (HBsAg) par une méthode comprenant deux traitements successifs avec le PEG 6000 a été décrite par Ph. ADAMOWICZ et al. (p. 37-49 INSERM SYMPOSIUM N° 18, HEPATITIS B VACCINE, Publ.

ELSEVIER, Amsterdam, Holland, 1981). Dans cette méthode de purification de l'HBsAg au départ de plasma, les complexes immuns et la plupart des lipoprotéines sont, dans une première étape, précipités du sérum par le PEG 6000 à une concentration de 5.5 % et, dans une seconde étape, l'HBsAg est précipité du surnageant isolé par addition de PEG à une concentration finale de 10 %.

En ce qui concerne la littérature sur les brevets,

-le brevet des Etats-Unis d'Amérique N° 3790552 décrit une méthode d'élimination de l'antigène associé à l'hépatite à partir d'une fraction protéinique et qui comprend une étape dans laquelle on utilise du PEG ayant un poids moléculaire de 200 à 6000 à raison d'une quantité de 12 à 30 % (p/v) pour précipiter ledit antigène.

-le brevet des Etats-Unis d'Amérique N° 3951937 décrit un procédé de purification de l'antigène de l'hépatite B (HBsAg) impliquant une double précipitation de l'HBsAg au moyen de PEG (4 -4,5 % en poids) ayant un poids moléculaire d'au moins 600.

-le brevet des Etats-Unis d'Amérique N° 3994870 décrit une méthode de purification de l'antigène de l'hépatite B (HBsAg) dans laquelle l'HBsAg est précipité par addition de 4 à 4,5 % en poids de PEG et est ensuite soumis à une chromatographie d'affinité à l'aide de concanavalline A comme adsorbant chromatographique.

-la demande de brevet européen publiée sous le numéro 0 112 506 décrit un procédé de production d'un vaccin préventif de l'infection à l'hépatite B à partir de plasma et qui comprend une précipitation au sulfate d'ammonium suivie d'adsorption sur du silicate colloïdal et de deux étapes de précipitations successives avec du PEG (ayant un poids moléculaire de 2000 à 10000) à raison de 3 à 7 % (p/v) de manière à précipiter le virus de l'hépatite B et les complexes immuns et à raison de 15 à 20 % (p/v) dans le surnageant de manière à précipiter l'HBsAg.

-Dans le domaine de l'isolement de l'alpha-1-antitrypsine, la demande de brevet japonais 59 129335 (résumé Derwent 84-217127) décrit la précipitation d'alpha-1-antitrypsine à partir d'une fraction plasmatique par addition d'une quantité de 15 à 20 % (p/v) de PEG.

Ainsi qu'il a été mentionné ci-dessus, une méthode de précipitation des lipoprotéines par application des interactions lipoprotéine-polysaccharide-métal a également été signalée précédemment - (par exemple : M. BURSTEIN et al. *Adv. Lip. Res.*, 11:67-108:1973 et A. VAN DALEN et al., *Biochim. Biophys. Acta* 147:421-427; 1967). Cette méthode est réalisée par interaction entre les lipoprotéines, un cation métallique bivalent et un polysaccharide acide et, dans ces conditions opératoires, la quantité de précipité est fonction de la concentration en cation métallique bivalent dans le milieu.

#### Description de l'invention

Le produit de départ utilisé dans le procédé de la présente invention (désigné également ici sous le nom d'"extrait brut") est le surnageant des cellules de levure produites génétiquement qui ont donné naissance à une protéine liée à la cellule et qui sont éclatées en présence d'un détergent non ionique, ainsi qu'il est bien connu dans la technique.

Conformément à la présente invention, on réalise alors la précipitation fractionnée par PEG - (désignée ci-après sous le nom de première étape) et, ensuite, soit une précipitation fractionnée par un cation métallique polyvalent (plus particulièrement un métal bivalent tel que le calcium ou le manganèse) soit un traitement au sulfate d'ammonium, ledit traitement au sulfate d'ammonium étant éventuellement précédé d'une ultrafiltration du surnageant provenant de la première étape.

La présente invention résulte de la découverte que, lorsqu'on utilise du PEG solide (p.ex. PEG 6000) à une concentration de 6 à 12 % (p/v) pour extraire l'HBsAg à partir de cellules de levure produites génétiquement et éclatées en présence d'un détergent non ionique, l'HBsAg ne précipite pas. De plus, par addition ultérieure de sulfate d'ammonium, il se crée un système biphasé qui se caractérise par le fait que l'HBsAg est présent dans la phase PEG tandis que la majeure partie des contaminants se trouve dans la phase aqueuse. Ce résultat est surprenant étant donné que ces conditions opératoires devraient provoquer la précipitation de l'HBsAg en vertu de ce que l'art antérieur nous apprend généralement au sujet d'autres milieux bruts. Le premier objectif de la présente invention est donc d'utiliser le PEG comme solvant pour isoler la protéine désirée produite par ces cultures de cellules de levure tandis que la majeure partie des contaminants sont éliminés par précipitation dans le milieu.

En ce qui concerne le poids moléculaire, le PEG existe soit sous forme solide, soit sous forme liquide, la limite entre les deux formes se situant autour d'un poids moléculaire de 1500.

Dans la première étape du procédé suivant la présente invention (qui pourrait être considérée également comme une étape de clarification), des PEGs ayant des poids moléculaires différents peuvent évidemment être utilisés en ajustant adéquatement leur concentration en fonction de leurs poids moléculaires respectifs.

Par exemple, la concentration en PEG varie, de préférence, de 6 à 12 % (p/v) lorsqu'on utilise du PEG solide (p.e. le PEG 6000) et de 10 à 35 % (v/v) - et de préférence de 20 à 30 % (v/v) lorsqu'on utilise du PEG liquide (p.e. le PEG 300 ou 400). Néanmoins, l'utilisation de PEG liquide est préférable pour des raisons techniques parmi lesquelles la facilité de la dernière ultrafiltration.

En conséquence, la présente invention décrit un procédé d'extraction et de purification de protéines, plus particulièrement de l'HBsAg ou de l'alpha-1-antitrypsine, au départ du surnageant de cellules de levure éclatées en présence d'un détergent non ionique (de préférence un détergent du type polysorbate), ledit surnageant étant désigné ici par "extrait brut", lequel procédé comprend une première étape dans laquelle on ajoute 6 à 12 % (p/v) de PEG solide ou de préférence 10 à 25 % (v/v) de PEG liquide à l'extrait brut amené à pH 6 ( $\pm 0,1$ ), les limites des concentrations en PEG susmentionnées étant principalement déterminées par le moment où la clarification est atteinte puisqu'une addition supplémentaire de PEG ne produit que des effets défavorables, c.à.d. une viscosité plus élevée du milieu et le risque d'une coprecipitation inopportune d'HBsAg ou d'alpha-1-antitrypsine.

Dans le procédé de la présente invention, le PEG est considéré comme un solvant organique polymérisé favorisant e.a. la précipitation des (lipo)-protéines dont le point isoélectrique se situe près du pH 6 et la solubilisation des autres (lipo)-protéines, c.à.d. celles qui présentent un point isoélectrique nettement différent et celles qui sont fortement hydrophobes.

La demanderesse a remarqué que l'étape de clarification par le PEG précipite environ 75 % des protéines contaminantes, 90 % des polysaccharides, 94 % des acides nucléiques et 45 % des lipides, tandis que c'est le contenu total en HBsAg ou en alpha-1-antitrypsine qui est pratiquement récupéré dans le surnageant.

Ainsi qu'il a été indiqué ci-dessus, la présente invention comprend une combinaison d'étapes dont la première est une étape de clarification en vue d'obtenir une solution partiellement purifiée, et éventuellement quelque peu opalescente de la protéine désirée.

Dans la deuxième étape du procédé de la présente invention, la solution obtenue et partiellement purifiée est traitée soit par un cation métallique polyvalent - plus particulièrement un cation métallique bivalent - comme une solution aqueuse de chlorure de calcium ou de manganèse à une concentration finale de 30 mM, soit par du sulfate d'ammonium, ce traitement au sulfate d'ammonium étant effectué soit après ultrafiltration de la solution opalescente par addition de sulfate d'ammonium solide jusqu'à 40 à 50 % de saturation, suivant la méthode classique dite de relargage pour précipiter la protéine désirée laquelle peut être reprise dans un tampon phosphate, soit par addition de sulfate d'ammonium au surnageant contenant le PEG jusqu'à 40 à 50 % de saturation pour former un système biphasé contenant la protéine désirée dans la phase PEG.

Chacune de ces variations dans la deuxième série d'étapes donne lieu à une solution limpide en protéine désirée effectivement purifiée en ce qui concerne son contenu en polysaccharides, acides nucléiques, lipides et protéines contaminantes.

La solution ainsi obtenue peut ensuite être traitée dans la troisième série d'étapes classiques telle que définie ci-dessus, p.ex. ultrafiltration éventuelle, filtration sur gel et chromatographie sur colonne et, dans un mode de réalisation préféré de la présente invention, cette troisième série d'étapes comprend successivement une ultrafiltration, une première filtration sur gel, une chromatographie sur colonne échangeuse d'anion faiblement alcalin dotée de groupes diéthylaminoéthyle (DEAE) et une seconde filtration sur gel ou centrifugation sur gradient de chlorure de césium de manière à obtenir une solution d'HBsAg ou d'alpha-1-antitrypsine hautement purifiée et appropriée à un usage médical.

Lorsqu'on teste le produit isolé après la seconde perméation sur gel par électrophorèse sur gel de polyacrylamide sur dodécyl sulfate de sodium (SDS PAGE), on constate qu'il est pur à plus de 98 % en ce qui concerne la bande 23 K qui caractérise l'HBsAg issu de la levure.

En conséquence le premier avantage du procédé de la présente invention réside dans le fait qu'il élimine effectivement tous les polysaccharides, les acides nucléiques, les lipides et les

matières protéiniques et un autre avantage de la présente invention est que la protéine désirée peut finalement être séparée avec un rendement relativement élevé.

Lorsque le procédé de la présente invention est appliqué à l'extrait brut d'HBsAg en provenance des cellules de levure procuites génétiquement, l'HBsAg obtenu est lié au détergent non ionique en formant avec celui-ci des micelles composites d'un diamètre d'environ 17 à 20 nm et la demanderesse a découvert que le rapport exprimant la quantité de détergent non ionique sur la quantité de protéines, de lipides totaux et de polysorbate dans la micelle composite de polysorbate varie de 15 à 35 % (p/v) lorsque les analyses sont effectuées par méthode colorimétrique pour les protéines (LOWRY), les lipides totaux (ZOLLNER) et le détergent non ionique - (HUDDLESTON et ALLRED).

Les micelles composites obtenues par le procédé de la présente invention à l'aide d'un détergent non ionique de type polysorbate constituent des composés nouveaux qui font également l'objet de la présente invention; ils sont immunogènes et peuvent être présentés dans une forme vaccinale comparable à l'HBsAg classique ainsi qu'il est bien connu dans la technique pour la préparation du vaccin à base du virus de l'hépatite B.

L'invention est illustrée par les exemples suivants qui ne limitent en rien la portée de l'invention.

#### Exemple 1

Un culot de cellules de levure (3850 g) provenant d'une souche produite génétiquement et exprimant l'HBsAg et qui se sont développées jusqu'à l'obtention de 30 g de cellules en poids sec par litre de culture sont mises en suspension dans 7,12 litres d'une solution de  $\text{Na}_2\text{HPO}_4$  (7,098 g/l).

Cette suspension est additionnée de 142,5 ml d'une solution de EDTA à 4 % (p/v), 38,5 ml de polysorbate 20 et 385 ml d'isopropanol contenant 2,7 g de fluorure de phénylméthylsulfonyl - (FPMS). Le pH est ajusté à 8,1 ( $\pm 0,1$ ) avec NaOH (10 % p/v dans l'eau). La suspension est refroidie dans un bain de glace et brisée par 2 passages à travers un homogénéiseur à billes de verre refroidi. L'homogénat (extrait brut) est ensuite centrifugé pendant 20 minutes à 13000 g.

On ajoute lentement, sous agitation, 2,5 l de PEG 400 à 7,7 l d'extrait brut que l'on maintient en dessous de 15° C. Le pH est ensuite ajusté à 6 (± 0,1) par addition d'acide acétique (5M) et le milieu est entreposé pendant une heure à 4° C avant d'être centrifugé à 7400-11000 g pendant 45 minutes.

Le surnageant est amené à pH 7 (± 0,05) avec NaOH N et on y ajoute 300 ml de CaCl<sub>2</sub> (c.à.d. 30 ml par litre de surnageant). Le pH est réajusté à 7 (± 0,05) avec NaOH N et entreposé à 4° C jusqu'au lendemain. La suspension est centrifugée à 3600 g pendant 45 minutes et le surnageant est ultrafiltré dans un appareil AMICON DC (appareil vendu par AMICON CORP., DANVERS, MA, USA) équipé d'une cartouche ayant un cut-off de 100000 daltons et, par ce moyen, il est d'abord concentré jusqu'à 1/4 (1500 ml) de son volume initial. La solution est ensuite lavée de façon continue avec 12,5 ml de trométamol 10mM ajusté à pH 8 (± 0,1) par addition d'acide chlorhydrique N (ce tampon est désigné ci-après par trométamol pH 8) et additionné de PMSF (1 mM), d'isopropanol (2,5 % v.v) et d'EDTA (2 mM) (concentrations finales). Le résidu est ensuite concentré ultérieurement jusqu'à 350 ml).

La solution concentrée est versée sur une colonne (diamètre 10 cm x 100 cm) contenant 7 l de FRACTOGEL<sup>®</sup> TSK HW65(F) (un gel semi-rigide composé de polymères vinyliques hydrophiles pourvus de nombreux groupes hydroxyyles à la surface de la matrice, présentant une dimension de particules de l'ordre de 32 à 63 µm, fabriqué et vendu par E. Merck, Darmstadt, RFA) équilibré dans un tampon trométamol HCl 10 mM pH 7 (± 0,01) additionné de 5 % (v.v) d'éthylène glycol.

Le pic contenant l'HBsAg est versé sur une colonne (diamètre 5 cm x 30 cm) de 300 ml de FRACTOGEL<sup>®</sup> TSK DEAE 650 (M) (un échangeur anionique faiblement basique dans lequel les groupes diéthylaminoéthyle sont liés aux groupes hydroxyle de la matrice du FRACTOGEL TSK HW65 par des liaisons étherées, fabriqué et vendu par E. Merck, Darmstadt, RFA) équilibré dans du trométamol pH 8 à 4° C. La colonne est lavée avec du trométamol pH 8 contenant NaCl (0,05 M) et l'HBsAg est élué avec NaCl (0,15 M) dans du trométamol pH 8.

L'éluat est appliqué sur une colonne de FRACTOGEL TSK HW65 (F) équilibrée avec un tampon Na<sub>2</sub>HPO<sub>4</sub>/NaH<sub>2</sub>PO<sub>4</sub> (10 mM) pH 6,8 additionné de NaCl (150 mM) pour donner une solution de micelles composites du type HBsAg/polysorbate.

Le taux de purification obtenu à chaque étape de la purification est indiqué au tableau I.

TABLEAU I

Fraction	Vol (l)	Total en HBsAg par RIE (mg)	Total en protéi- nes (g)	Total en polysac- charides (g)	Total en acides nucléi- ques (mg)	Total en lipides (g)
Extrait brut	7,7	1232	259	114	32500	104
Surna- geant PEG	8,02	1100	64	9,5	641	53,7
Surna- geant CaCl <sub>2</sub>	8	1190	37	3,8	512	27,2
Rétentat Pic	.350	1320	26,7	.744	37,6	3,596
HBsAg TSK	1,06	736	.935	.014	16	.352
Eluat TSK-DEAE	.15	722	.428	.0084	2,6	.275
Pic HBsAg TSK	.45	1117	.244	.0094	2	.179

RI3 : Radio-immuno essai

Le tableau II suivant résumé sommairement la composition des micelles composites de 3 lots différents de vaccins obtenus par le procédé ci-dessus avec du polysorbate 20.

45

50

55

7

TABLEAU II

Lot	Protéines	Total en	Polysorbate 20	Rapport
	( $\mu\text{g/ml}$ ) (A)	lipides ( $\mu\text{g/ml}$ ) (B)	( $\mu\text{g/ml}$ ) (C)	$\frac{C}{A/B/C}$
I	20	20	8,5	18
II	20	16	8,4	19
III	20	14,5	15,6	31

## Exemple 2

On ajoute lentement 882 ml de PEG 400 à 2.650 l d'extrait brut préparé comme il est décrit à l'exemple 1 et on ajuste le pH à 6 ( $\pm 0.1$ ). Le milieu est maintenu à 4° C pendant une heure et centrifugé ensuite à 7400 g. Le surnageant est concentré à 1000 ml sur un appareil AMICON DC équipé d'une cartouche ayant un cut-off de 100000 daltons et lavé ensuite avec 3 volumes de trométamol (20 mM) pH 8. Sous agitation et à 4° C. on ajoute ensuite lentement 277 g de sulfate d'ammonium en poudre au rétentat. Après 1 heure à 4° C. le précipité est centrifugé pendant 15 min. à 1000 g. Le surnageant est écarté et le précipité est dissout dans 400 ml de trométamol (20 mM) pH 8 additionné de 1 mM de PMSF. La solution est ultrafiltrée sur un appareil AMICON DC équipé d'une cartouche ayant un cut-off de 10<sup>5</sup> daltons. Le

rétenant (500 ml) est lavé avec 2 volumes de trométamol pH 8 et appliqué ensuite sur une colonne contenant 300 ml de FRACTOGEL TSK DEAE 650M équilibré avec du trométamol pH 8. Lorsque le passage de l'échantillon est achevé, la colonne est lavée avec NaCl 0.05 M dans du trométamol pH 8 et un gradient linéaire en NaCl (0.05 M -0.5 M) est ensuite appliqué sur la colonne. La fraction éluée avec le NaCl 0.15 M contient l'HBsAg; elle est concentrée à 50 ml dans un appareil AMICON DC équipé d'une cartouche ayant un cut-off de 10000 daltons et appliquée sur une colonne - (diamètre 50 x 100 cm) de Sepharose 4B-Cl (un gel d'agarose fabriqué et vendu par Pharmacia Fine Chemicals, Uppsala, Suède) équilibrée avec du trométamol (20 mM) pH 8 additionné de 5 % d'éthylène glycol pour donner une solution de micelles composites de type HBsAg/polysorbate.

Le taux de purification obtenu à chaque étape est indiqué au tableau III.

40

45

50

55

8



TABLEAU III

Fraction	Vol	Total en HBsAg par RIE (ml)	Total en protéi- nes (g)	Total en polysac- charides (g)	Total en acides nucléi- ques (mg)
Extrait brut	2650	188	114,721	36,9	9250
Surnageant PEG	2530	160	22,755	3,321	536
Rétentat 10 <sup>5</sup>	1000	150	7,468	.181	175
AMS-culot	400	80,8	1,248	.0248	160
Rétentat 10 <sup>6</sup>	1000	72,4	.529	.029	88
Eluat TSK-DEAE	560	44	.157	<0,073	3
Pic HBsAg					
Sépharose 4B-CL	200	23	.030	.0003	.05

RIE : Radio-immuno essai

## Exemple 3

On clarifie 2 litres d'extrait brut d'HBsAg avec du PEG 400 comme il est décrit à l'exemple 1. On y ajoute 450 g de sulfate d'ammonium (SAM) en poudre sous agitation (la concentration finale correspond à une saturation à 45 % en SAM. Après solubilisation du SAM, la solution est conservée à 4° C pendant une période de 3 heures au terme de laquelle on obtient deux phases distinctes que l'on sépare dans une ampoule à décantation : une phase supérieure limpide (jaune) (phase PEG 400)

et une phase inférieure limpide (phase aqueuse), chacune des phases représentant environ ± 50 % du volume original. La phase aqueuse est éliminée et la phase supérieure est ultrafiltrée sur un appareil AMICON DC, comme il est décrit pour le surnageant du CaCl<sub>2</sub> à l'exemple 1. Le processus de purification est ensuite poursuivi ultérieurement comme il est décrit à l'exemple 1 pour donner une solution de micelles composites de type HBsAg/polysorbate.

Le taux de purification obtenu à chaque étape est indiqué au tableau IV.

TABLEAU IV

Fraction	Vol	Total	Total	Total	Total
		en	en	en	en
		HBsAg	protéi-	polysac-	acides
		par	nes	charides	nucléi-
		RIE			ques
	(ml)	(mg)	(g)	(g)	(mg)
Extrait brut	2000	99	55	22	5588
Surnageant PEG	2000	92	13,2	5,68	243
phase PEG	1000	98	2,8	1,536	98
Retentat	50	98	1,5	.14	6,66
Pic HBsAg TSK	123	52	.288	.0077	2,84
Eluat TSK-DEAE	25	54	.0377	.0011	.44
Pic HBsAg TSK	60	35	.0207	.00154	.39

RIE : Radio-immuno essai

#### Exemple 4

On clarifie 2 litres d'extrait brut en HBsAg avec du PEG 400 comme il est décrit à l'exemple 1. On y ajoute 450 g de sulfate d'ammonium (SAM) en poudre sous agitation (la concentration finale correspond à une saturation à 45 % en SAM). Après solubilisation du SAM, la solution est conservée à 4° C pendant une période de 3 heures au terme de laquelle on obtient deux phases distinctes que l'on sépare dans une ampoule à décantation : une phase supérieure limpide (jaune) (phase PEG 400) et une phase inférieure limpide (phase aqueuse), chacune des phases représentant environ ± 50 % du volume original. La phase aqueuse est éliminée et une partie aliquote d'un litre de la phase PEG - (phase supérieure) est diluée deux fois avec du

trométamol 10 mM à pH 8 et appliquée sur une colonne contenant 300 ml de gel du type FRAC-TOGEL TSK-DEAE 650 (M) équilibrée avec du trométamol à pH 8, à raison d'un débit de 100 ml par heure. Le gel est lavé avec du trométamol à pH 8 contenant NaCl 0,05 M et l'HBsAg est ensuite élué avec un gradient en NaCl (NaCl 0,05 à 0,5 M dans le trométamol à pH 8). Un pic HBsAg est élué avec NaCl 0,15 M et après concentration dans un appareil AMICON DC équipé d'une cartouche ayant un cut-off de 10000 daltons, le rétentat est appliqué sur une colonne (diamètre 5 x 100 cm) de Sépharose 4B-Cl équilibrée avec du trométamol pH 8 contenant 5 % (v/v) d'éthylène glycol pour donner un pic de micelles composites de type HBsAg/polysorbate/

Le taux de purification obtenu à chaque étape est indiqué au tableau V.

50

55

10

TABLEAU V

Fraction	Vol	Total	Total	Total	Total
		en	en	en	en
		HBsAg	protéi-	polysac-	acides
		par	nes	charides	nucléi-
		RIE			ques
	(ml)	(mg)	(g)	(g)	(mg)
Extrait brut	2100	200	60,600	19,000	4860
phase PEG	1200	210	3,100	1,436	80
Eluat TSK-DEAE	460	134	.750	.0048	05
Pic HBsAg/					
Sépharose 4B-CL	150	110	.120	.0014	0.3

RIE : Radio-immuno essai

## Exemple 5

On clarifie 500 ml d'extrait brut en HBsAg par addition lente d'une solution de 50 g de PEG 6000 dans 160 ml d'eau à 4° C, sous agitation. On ajuste le pH à 6,1 par addition d'acide acétique 5M. Après une heure de stockage à 4° C, l'extrait est centrifugé pendant 15 min. à 7000 g. On ajoute ensuite, sous agitation, 157 g de sulfate d'ammonium en poudre au surnageant (la concentration finale correspond à 50 % de sulfate d'ammonium saturé). Après solubilisation, la solution est conservée à 4° C pendant une période de 3 heures au terme de laquelle on obtient deux phases distinctes que l'on sépare dans une ampoule à décantation. La phase supérieure (phase PEG con-

25

30

35

40

45

50

55

11

tenant l'HBsAg) est diluée deux fois avec du trométamol à pH 8 et appliquée ensuite sur une colonne contenant 300 ml de gel de TSK-DEAE 650 (M) équilibrée avec du trométamol à pH 8 à raison d'un débit de 100 ml par heure. Le gel est lavé avec du trométamol à pH 8 contenant NaCl 0,15 M, concentré dans un appareil AMICON DC équipé d'une cartouche ayant un cut-off de 10000 daltons et le rétentat est appliqué ensuite sur une colonne (diamètre 5 x 100 cm) de 2 litres de FRACTOGEL TSK-HW65(F) équilibré avec du trométamol à pH 8 contenant 10 % (v/v) d'éthylène glycol pour donner une solution de micelles composites de type HBsAg/polysorbate.

Le taux de purification à chaque étape est indiqué au tableau VI.

TABLEAU VII

Fraction	Vol (ml)	Total en AAT par ELISA (mg)	Total en protéi- nes (g)	Total en polysac- charides (g)
Extrait brut	120	180	4.018	.538
Surnageant PEG	100	212	1.300	.290
Surnageant $\text{CaCl}_2$	100	194	.660	.110
Eluat TSK-DEAE	520	180	.210	.0052

AAT : alpha-1-antitrypsine

## Exemple 7

La technique est telle que décrite à l'exemple 1, mais on remplace les 300 ml de  $\text{CaCl}_2$  M qui y sont spécifiés par 300 ml de  $\text{MnCl}_2$  M. Les caractéristiques du produit final sont analogues à celles du produit obtenu à la fin de l'exemple 1.

## Exemple 8

La technique est telle que décrite à l'exemple 1, mais on remplace les 38,5 ml de polysorbate 20 qui y sont spécifiés par 20 ml de Triton X-100 (un produit fabriqué par Rohm et Haas; Darmstadt, RFA). Les caractéristiques du produit final sont analogues à celles du produit obtenu à la fin de l'exemple 1.

## Exemple 9

La technique est telle que décrite à l'exemple 1, mais on remplace les 38,5 ml de polysorbate 20 qui y sont spécifiés par 38,5 ml de polysorbate 80. Les caractéristiques du produit final sont analogues à celles du produit obtenu à la fin de l'exemple 1.

## Exemple 10

La solution du micelle composite de type HB-sAg obtenu à la fin de l'exemple 1 est ajustée à un contenu protéinique de 10  $\mu\text{g}$  par millilitre par addition de NaCl, de tampon phosphate ( $\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$ ) et d'ALHYDROGEL<sup>R</sup> (un gel d'hydroxyde d'aluminium fabriqué et vendu par

SUPERHOS Export Co. Copenhagen, Danemark) jusqu'à des concentrations finales respectives de 0,9 % (p/v), 20 mM et 0,15 % (p/v) en  $\text{Al}(\text{OH})_3$ , le pH final étant de 6,9. La préparation est stérilisée et répartie dans des fioles en verre de 2 ml contenant chacune une dose unitaire d'un ml de vaccin.

## Exemple 11

Les doses unitaires du vaccin de l'exemple 10 ont été administrées par voie intramusculaire à deux séries de sujets séronégatifs à raison de 2 injections successives, à un mois d'intervalle. Bien que ces injections devraient être considérées comme une administration de sensibilisation qui devrait être suivie d'un rappel, p.ex. 2 mois après la première injection, une réponse immunitaire positive a déjà été observée respectivement un et deux mois après la première administration.

Dans la première série (comprenant 32 sujets séronégatifs), un mois après la première administration, le taux de séroconversion était de 20/32, c'est-à-dire de 62,5 % avec un titre géométrique moyen (TGM) de 9,95 unités milli-internationales (UMI) et un mois après la seconde administration, le taux de séroconversion était de 31/32, c'est-à-dire de 96,87 % avec un TGM de 36 UMI.

Dans la seconde série (comprenant 46 sujets séronégatifs), un mois après la première administration, le taux de séroconversion était de 31/46, c'est-à-dire de 67,4 % avec un TGM de 13,6 UMI.

L'enregistrement des signes et symptômes cliniques chez les vaccinés n'a révélé aucune élévation de température et aucune réaction à l'endroit de l'administration.

#### Revendications

Revendication pour les pays contractants : UE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Procédé d'extraction et de purification d'une protéine liée à la cellule à partir du surnageant d'une culture de cellules de levure manipulées génétiquement ayant produit ladite protéine et éclatées en présence d'un détergent non ionique, caractérisé en ce qu'on ajuste le pH du surnageant à 6 ( $\pm 0,1$ ), on y ajoute du polyéthylène glycol liquide ou solide jusqu'à clarification dudit surnageant et on traite le surnageant clarifié soit par un cation métallique bivalent, soit, après ultrafiltration éventuelle, par du sulfate d'ammonium pour séparer ladite protéine.

2. Procédé suivant la revendication 1, dans lequel on ajoute du polyéthylène glycol liquide à une concentration finale comprise entre 10 et 35 % (v/v).

3. Procédé suivant la revendication 2, dans lequel on ajoute du polyéthylène glycol solide à une concentration finale comprise entre 6 et 12 % (p/v).

4. Procédé suivant l'une quelconque des revendications 1 à 3, dans lequel la protéine est l'antigène de surface de l'hépatite B.

5. Procédé suivant l'une quelconque des revendications 1 à 3, dans lequel la protéine est l' $\alpha$ -antitrypsine.

6. Procédé suivant l'une quelconque des revendications 1 à 5, dans lequel le surnageant clarifié est traité par un cation métallique bivalent.

7. Procédé suivant la revendication 6, dans lequel le cation métallique bivalent est le cation calcium ou manganèse.

8. Procédé suivant l'une quelconque des revendications 6 et 7, qui comprend en outre la séparation du précipité, l'ultrafiltration de la solution, la perméation sur gel du rétentat, la chromatographie sur échangeurs d'ions du pic contenant la protéine et éventuellement une seconde perméation sur gel ou une centrifugation sur gradient de chlorure de

cesium.

9. Procédé suivant l'une quelconque des revendications 1 à 5, dans lequel on traite le surnageant clarifié par du sulfate d'ammonium jusqu'à 40 à 50 % de saturation et on isole la protéine purifiée dans la phase polyéthylène glycol.

10. Procédé suivant la revendication 9, dans lequel la phase polyéthylène glycol est soumise en outre à une éventuelle ultrafiltration suivie d'une perméation sur gel et d'une chromatographie sur échangeurs d'ions.

11. Procédé suivant la revendication 9, dans lequel la phase polyéthylène glycol est soumise en outre à une éventuelle ultrafiltration suivie d'une chromatographie sur échangeurs d'ions et d'une perméation sur gel.

12. Procédé suivant la revendication 2, dans lequel le surnageant clarifié est soumis à une ultrafiltration et le rétentat est traité par du sulfate d'ammonium jusqu'à 40 à 50 % de saturation afin de précipiter la protéine qui est reprise dans un tampon adéquat.

13. Procédé suivant la revendication 12 qui comprend en outre l'ultrafiltration, la perméation sur gel et la chromatographie sur échangeurs d'ions.

14. Procédé suivant l'une quelconque des revendications 1 à 5, dans lequel le détergent non ionique est un polysorbate.

15. Procédé suivant la revendication 14, dans lequel le polysorbate est le polysorbate 20.

16. Micelle composite à base d'antigène de surface de l'hépatite B et de polysorbate contenant de 15 à 35 % (p/p) de polysorbate.

17. Vaccin contre l'hépatite B contenant à titre d'ingrédient actif une micelle composite suivant la revendication 16.

#### Revendications pour l'Etat Contractant AT

18. Procédé d'extraction et de purification d'une protéine liée à la cellule à partir du surnageant d'une culture de cellules de levure manipulées génétiquement ayant produit ladite protéine et éclatées en présence d'un détergent non ionique, caractérisé en ce qu'on ajuste le pH du surnageant à 6 ( $\pm 0,1$ ), on y ajoute du polyéthylène glycol liquide ou solide jusqu'à clarification dudit surnageant

geant et on traite le surnageant clarifié soit par un cation métallique bivalent, soit, après ultrafiltration éventuelle, par du sulfate d'ammonium pour séparer ladite protéine.

2. Procédé suivant la revendication 1, dans lequel on ajoute du polyéthylène glycol liquide à une concentration finale comprise entre 10 et 35 % (v/v).

3. Procédé suivant la revendication 2, dans lequel on ajoute du polyéthylène glycol solide à une concentration finale comprise entre 6 et 12 % (p/v).

4. Procédé suivant l'une quelconque des revendications 1 à 3, dans lequel la protéine est l'antigène de surface de l'hépatite B.

5. Procédé suivant l'une quelconque des revendications 1 à 3, dans lequel la protéine est l'alpha-1-antitrypsine.

6. Procédé suivant l'une quelconque des revendications 1 à 5, dans lequel le surnageant clarifié est traité par un cation métallique bivalent.

7. Procédé suivant la revendication 6, dans lequel le cation métallique bivalent est le cation calcium ou manganèse.

8. Procédé suivant l'une quelconque des revendications 6 et 7, qui comprend en outre la séparation du précipité, l'ultrafiltration de la solution, la perméation sur gel du rétentat, la chromatographie sur échangeurs d'ions du pic contenant la protéine et éventuellement une seconde perméation sur gel ou une centrifugation sur gradient de chlorure de

cesium.

9. Procédé suivant l'une quelconque des revendications 1 à 5, dans lequel on traite le surnageant clarifié par du sulfate d'ammonium jusqu'à 40 à 50 % de saturation et on isole la protéine purifiée dans la phase polyéthylène glycol.

10. Procédé suivant la revendication 9, dans lequel la phase polyéthylène glycol est soumise en outre à une éventuelle ultrafiltration suivie d'une perméation sur gel et d'une chromatographie sur échangeurs d'ions.

11. Procédé suivant la revendication 9, dans lequel la phase polyéthylène glycol est soumise en outre à une éventuelle ultrafiltration suivie d'une chromatographie sur échangeurs d'ions et d'une perméation sur gel.

12. Procédé suivant la revendication 2, dans lequel le surnageant clarifié est soumis à une ultrafiltration et le rétentat est traité par du sulfate d'ammonium jusqu'à 40 à 50 % de saturation afin de précipiter la protéine qui est reprise dans un tampon adéquat.

13. Procédé suivant la revendication 12 qui comprend en outre l'ultrafiltration, la perméation sur gel et la chromatographie sur échangeurs d'ions.

14. Procédé suivant l'une quelconque des revendications 1 à 5, dans lequel le détergent non ionique est un polysorbate.

15. Procédé suivant la revendication 14, dans lequel le polysorbate est le polysorbate 20.

40

45

50

55

15

DERWENT-ACC-NO: 1990-192954

DERWENT-WEEK: 199025

COPYRIGHT 2005 DERWENT INFORMATION LTD

TITLE: Solid unit dosage form of piperidino-alkanol cpds. - contg. surfactant and carbonate salt for ready bio-availability of antihistamine, broncho dilator or anti-allergy agent

INVENTOR: DOMET, J; SHAH, D N

PRIORITY-DATA: 1989US-0325254 (March 17, 1989), 1987US-0105928 (October 7, 1987), 1987US-0117166 (November 4, 1987), 1988US-0152689 (February 5, 1988)

Search Selected

Search ALL

Clear

## PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
<input type="checkbox"/> <u>US 4929605 A</u>	May 29, 1990		000	

INT-CL (IPC): A61K 9/20; A61K 31/44; A61K 47/00

ABSTRACTED-PUB-NO: DE 3873011G

## BASIC-ABSTRACT:

A pharmaceutical compsn. in solid unit dosage form comprises (a) a therapeutically effective amt. of a piperidino alkanol cpd. or one of its salts, (b) a non-ionic or cationic surfactant in an amt. of 0.1-6 wt.% of the compsn., and (c) a carbonate salt in an amt. of 2-50 wt.% of the compsn.

Pref. cpds. (a) include alpha-(4-(1,1-dimethylethyl)phenyl) -4-hydroxydiphenylmethyl -1-piperidine butanol, which may be present, e.g. in an amt. of 10-120mg, esp. 60 or 120 mg. The pref. carbonate is calcium carbonate, used in amt. of 2-25 wt.%, more pref. 12-18 wt.% and esp. 15 wt.%. Non-ionic surfactant is pref. present in an amt. of 2-4 wt.%, esp. 3 wt.% and the pref. surfactant is polysorbate 80. There may also be present microcrystalline cellulose (35 wt.%), pregelatinised corn starch (30 wt.%), sodium starch glycolate (5 wt.%) and magnesium stearate (0.5 wt.%). A suitable cationic surfactant is cetyl pyridinium chloride.

USE/ADVANTAGE - Cpds. (a) are antihistamines, antiallergy agents and bronchodilators. The present compsns. contg. them allow efficient and immediate absorption and bioavailability of the cpds. after their oral admin.

ABSTRACTED-PUB-NO:

## EP 310999B EQUIVALENT-ABSTRACTS:

Pharmaceutical compsn. in solid unit dosage form comprises (1) at least one piperidino-alkanol cpd. (I), or its pharmaceutically acceptable salt; (2) at least one surfactant at 0.1-6 wt.% and (3) 2-50 wt.% CaCO<sub>3</sub>. Specifically, (I) is alpha-(4-(1,1-dimethylethyl) phenyl-4-(hydroxydiphen-yl methyl)- 1-piperidino-butanol (Ia); the surfactant is nonionic or cationic (pref. 'Polysorbate 80', polyethylene glycol (PEG) or cetylpyridinium chloride) at 0.5-4, esp. 1-3 wt.% and CaCO<sub>3</sub> is 2-25, esp. 12-15, wt.%.

USE/ADVANTAGE - (I) are variously useful as antihistamines, antiallergic agents or bronchodilators, particularly (Ia) is used to treat seasonal allergic rhinitis. Although (I) have very low solubility in water, in these compsns. they provide efficient and immediate absorption and bioavailability, following oral

admin. Pref. unit doses contain about 60 or 120 mg (I).

A pharmaceutical composition in solid unit dosage form comprising a) a therapeutically effective amount of at least one piperidinoalkanol compound of the formula (1), wherein R1 is hydrogen or hydroxy; R2 is hydrogen; or R1 and R2 taken together form a second bond between the carbon atoms bearing R1 and R2; n is an integer of from 1 to 5; in case n is an integer of from 1 to 3, Z is thienyl, phenyl or substituted phenyl wherein the substituents on the substituted phenyl may be attached at the ortho, meta, or para positions of the substituted phenyl ring and are selected from a halogen atom, a straight or branched lower alkyl chain of from 1 to 4 carbon atoms, a lower alkoxy group of from 1 to 4 carbon atoms, a di(lower alkyl)amino group, or a saturated monocyclic heterocyclic ring selected from pyrrolidino, piperidino, morpholino, or N-(lower alkyl)-piperizino; or in case n is an integer of from 1 to 5, Z is a gp. of formula (i), wherein R3 is -CH<sub>3</sub>, -CH<sub>2</sub>OH, -COOH or -COOalkyl wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched; each of A and B is hydrogen or hydroxy; with the provisos that at least one of A or B is hydrogen and one of A or B is other than hydrogen when R3 is -CH<sub>3</sub>; and pharmaceutically acceptable salts and individual optical isomers thereof, b) one or more nonionic surfactants in an amount of from 0.1% to 6% by weight of the composition, c) calcium carbonate in an amount of from 2% to 50% by weight of the composition.

US 4929605A



First Hit

L8: Entry 34 of 160

File: PGPB

Dec 25, 2003

DOCUMENT-IDENTIFIER: US 20030235595 A1

TITLE: Oil-containing, orally administrable pharmaceutical composition for improved delivery of a therapeutic agent

## CLAIMS:

22. The composition of claim 21, wherein the at least one hydrophilic surfactant is selected from polysorbate 80, PEG-35 castor oil, and PEG-40 castor oil.

76. A pharmaceutical composition comprising: (a) a carrier comprising glyceryl tricaprilate/caprate, at least one hydrophilic surfactant selected from the group consisting of tocopheryl PEG-1000 succinate, polysorbate 80, PEG-35 castor oil, PEG-40 hydrogenated castor oil, PEG-8 caprylic/capric glycerides, lauroyl macrogol-32 glycerides, stearyl macrogol glyceride, and mixtures thereof; and at least one hydrophobic surfactant selected from the group consisting of glyceryl caprylate, glyceryl caprylate/caprate and mixtures thereof; and (b) a therapeutically effective amount of fenofibrate, wherein the triglyceride and surfactants are present in amounts that are pharmaceutically acceptable and selected so that upon admixture of the composition with an aqueous solution in an aqueous solution to composition ratio of about 100:1 by weight, a clear aqueous dispersion having an absorbance of less than about 0.3 at 400 nm is provided.

77. The composition of claim 76, wherein the at least one hydrophilic surfactant is selected from the group consisting of tocopheryl PEG-1000 succinate, polysorbate 80, and mixtures thereof.

First Hit

L8: Entry 34 of 160

File: PGPB

Dec 25, 2003

PGPUB-DOCUMENT-NUMBER: 20030235595  
PGPUB-FILING-TYPE: new  
DOCUMENT-IDENTIFIER: US 20030235595 A1

TITLE: Oil-containing, orally administrable pharmaceutical composition for improved delivery of a therapeutic agent

PUBLICATION-DATE: December 25, 2003

## INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Chen, Feng-Jing	Salt Lake City	UT	US	
Patel, Mahesh V.	Salt Lake City	UT	US	

APPL-NO: 10/ 397969 [PALM]  
DATE FILED: March 25, 2003

## RELATED-US-APPL-DATA:

Application 10/397969 is a continuation-in-part-of US application 09/877541, filed June 8, 2001, PENDING  
Application 09/877541 is a continuation-in-part-of US application 09/345615, filed June 30, 1999, US Patent No. 6267985  
Application 09/877541 is a continuation-in-part-of US application 09/751968, filed December 29, 2000, US Patent No. 6458383  
Application 09/751968 is a continuation-in-part-of US application 09/375636, filed August 17, 1999, US Patent No. 6309663

INT-CL: [07] A61 K 9/00, A61 K 31/192

US-CL-PUBLISHED: 424/400; 514/571  
US-CL-CURRENT: 424/400; 514/571

## ABSTRACT:

The present invention relates to oral pharmaceutical compositions and methods for improved delivery of therapeutic agents, e.g., lipid-regulating agents. Compositions of the present invention include a carrier, where the carrier contains a combination of a triglyceride and at least two surfactants, at least one of which is hydrophilic. Upon dilution with an aqueous medium, the composition forms a clear, aqueous dispersion. The invention also pertains to methods for treating lipid disorders such as hypercholesterolemia, hypertriglyceridemia, and mixed dyslipidemia by oral administration of the compositions provided.

## CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation-in-part of U.S. patent application Ser. No. 09/877,541, filed Jun. 8, 2001, which is a continuation-in-part of U.S. patent application Ser. No. 09/345,615, filed Jun. 30, 1999, issued on Jul. 31, 2001 as



stable or free sulfhydryl-containing protein formulation, and a nonionic surfactant.

The method of using the sustained release composition of the present invention comprises providing a therapeutically effective blood level of biologically active acid-stable or free sulfhydryl-containing protein in a subject for a sustained period by administering to the subject a dose of the sustained release composition described herein.

#### RELATED APPLICATION

[0001] This application is a continuation of U.S. application Ser. No. 09/501,934, filed Feb. 10, 2000. The entire teachings of the above application are incorporated herein by reference.

DOCUMENT-IDENTIFIER: US 20030138491 A1

TITLE: Microencapsulation and sustained release of biologically active acid-stable or free sulfhydryl-containing proteins

CLAIMS:

10. The sustained release composition of claim 1 wherein the surfactant is selected from the group consisting of: poloxamers, polysorbates, polyethyleneglycols, polyoxyethylene fatty acid esters, bile salts, benzalkonium chloride, polyoxyethylene (40) monostearate and combinations thereof.

21. The sustained release composition of claim 20 wherein the nonionic surfactant is selected from the group consisting of: poloxamers, polysorbates, polyethyleneglycol, polyoxyethylene fatty acid esters and combinations thereof.

DOCUMENT-IDENTIFIER: US 20030064030 A1

TITLE: Ultrasound contrast agents and methods of making and using them

CLAIMS:

75. A vesicular composition according to claim 74 wherein said polymer is selected from the group consisting of poly(ethylene glycol) (PEG), poly(vinylpyrrolidone), polyoxomers, polysorbate and polyvinyl alcohol.

PGPUB-DOCUMENT-NUMBER: 20030064030  
PGPUB-FILING-TYPE: new  
DOCUMENT-IDENTIFIER: US 20030064030 A1

TITLE: Ultrasound contrast agents and methods of making and using them

PUBLICATION-DATE: April 3, 2003

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Schneider, Michel	Troinex		CH	
Yan, Feng	Carouge		CH	
Garcel, Nadine	Ambilly		FR	
Grenier, Pascal	Ambilly		FR	
Puginier, Jerome	Le Chable-Beaumont		FR	
Barrau, Marie-Bernadette	Geneve		CH	
Bussat, Philippe	Feigeres		FR	
hybl, Eva	Heidelberg		DE	
Bichon, Daniel	Montpellier		FR	

APPL-NO: 10/ 102684 [PALM]

DATE FILED: March 22, 2002

RELATED-US-APPL-DATA:

Application 10/102684 is a continuation-of US application 09/225293, filed January 5, 1999,  
ABANDONED

Application 09/225293 is a continuation-in-part-of US application 08/033435, filed March 18, 1993,  
ABANDONED

Application 08/033435 is a division-of US application 07/695343, filed May 3, 1991, ABANDONED

Application 10/102684 is a continuation-of US application 09/225293, filed January 5, 1999,  
ABANDONED

Application 09/225293 is a continuation-in-part-of US application 08/853936, filed May 9, 1997, US  
Patent No. 6110443

Application 08/853936 is a division-of US application 08/456385, filed June 1, 1995, US Patent No.  
5658551

Application 08/456385 is a division-of US application 08/315347, filed September 30, 1994, US Patent  
No. 5531980

Application 08/315347 is a division-of US application 08/128540, filed September 29, 1993, US Patent  
No. 5380519

Application 08/128540 is a division-of US application 07/775989, filed November 20, 1991, US Patent  
No. 5271928

Application 10/102684 is a continuation-of US application 09/225293, filed January 5, 1999,  
ABANDONED

Application 09/225293 is a continuation-in-part-of US application 08/740653, filed October 31, 1996,  
PENDING

Application 08/740653 is a division-of US application 08/350588, filed December 6, 1994, US Patent  
No. 5518991

Application 08/350588 is a division-of US application 07/991237, filed December 16, 1992, US Patent

No. 5413774

## FOREIGN-APPL-PRIORITY-DATA:

COUNTRY	APPL-NO	DOC-ID	APPL-DATE
EP	90810262.7	1990EP-90810262.7	April 2, 1990
EP	90810367.4	1990EP-90810367.4	May 18, 1990
WO	PCT/EP91/00620	1991WO-PCT/EP91/00620	April 2, 1991
EP	92810046.0	1992EP-92810046.0	January 23, 1992

INT-CL: [07] A61 L 9/04

US-CL-PUBLISHED: 424/45

US-CL-CURRENT: 424/45

## ABSTRACT:

Gas or air filled microbubble suspensions in aqueous phases usable as imaging contrast agents in ultrasonic echography. They contain laminarized surfactants and, optionally, hydrophilic stabilizers. The laminarized surfactants can be in the form of liposomes. The suspensions are obtained by exposing the laminarized surfactants to air or a gas before or after admixing with an aqueous phase. One can impart outstanding resistance against collapse under pressure to these gas-filled microbubbles used as contrast agents in ultrasonic echography by using as fillers gases whose solubility in water, expressed in liter of gas by liter of water under standard conditions, divided by the square root of the molecular weight does not exceed 0.003.

[0001] This application is a continuation-in-part of Ser. No. 08/033,435, filed Mar. 18, 1993, which is a divisional of Ser. No. 07/695,343, filed May 3, 1991, which originated from EP 90810367.4, filed May 18, 1990. This application is also a continuation-in-part of Ser. No. 08/853,936, filed May 9, 1997, which is still pending, which is a divisional of Ser. No. 08/456,385, filed Jun. 1, 1995, now U.S. Pat. No. 5,658,551, which is a divisional of Ser. No. 08/315,347, filed Sep. 30, 1994, now U.S. Pat. No. 5,531,980, which is a divisional of Ser. No. 08/128,540, filed Sep. 29, 1993, now U.S. Pat. No. 5,380,519, which is a divisional of Ser. No. 07/775,989, filed Nov. 20, 1991, now U.S. Pat. No. 5,271,928, which originated from PCT/EP91/00620, filed Apr. 2, 1991, and EP 90810262.7, filed Apr. 2, 1990. This application is also a continuation-in-part of Ser. No. 08/740,653, filed Oct. 31, 1996, which is still pending, which is a divisional of Ser. No. 08/350,588, filed Jan. 30, 1995, now U.S. Pat. No. 5,578,292, which is a divisional of Ser. No. 07/991,237, filed Dec. 16, 1992, now U.S. Pat. No. 5,413,774, which originated from EP 92810046.0, filed Jan. 24, 1992. All of the aforementioned applications are hereby incorporated by reference herein in their entirety.



SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1951-2005/Jun W3  
(c) format only 2005 The Dialog Corp.  
File 5:Biosis Previews(R) 1969-2005/Jun W3  
(c) 2005 BIOSIS  
File 73:EMBASE 1974-2005/Jun 21  
(c) 2005 Elsevier Science B.V.  
File 286:Biotechnology Directory Current May B2  
(c) 2005 BioCommerce Data Ltd.  
File 10:AGRICOLA 70-2005/Jun  
(c) format only 2005 The Dialog Corporation  
File 156:ToxFile 1965-2005/Jun W3  
(c) format only 2005 The Dialog Corporation  
\*File 156: ToxFile has been reloaded with the 2005 MeSH.  
Please see HELP NEWS 156 for details.  
File 357:Derwent Biotech Res. 1982-2005/Jun W3  
(c) 2005 Thomson Derwent & ISI  
File 144:Pascal 1973-2005/Jun W2  
(c) 2005 INIST/CNRS  
File 160:Gale Group PROMT(R) 1972-1989  
(c) 1999 The Gale Group  
File 124:CLAIMS/REFERENCE 2001/2004Q4 (c) 2005 IFI/CLAIMS(R) PATENT SERVICES  
File 654:US Pat.Full. 1976-2005/Jun 21  
(c) Format only 2005 The Dialog Corp.  
File 347:JAPIO Nov 1976-2005/Feb(Updated 050606)  
(c) 2005 JPO & JAPIO  
File 16:Gale Group PROMT(R) 1990-2005/Jun 23  
(c) 2005 The Gale Group  
File 185:Zoological Record Online(R) 1978-2005/Jul  
(c) 2005 BIOSIS  
File 340:CLAIMS(R)/US Patent 1950-05/Jun 21  
(c) 2005 IFI/CLAIMS(R)  
File 94:JICST-EPlus 1985-2005/May W1  
(c)2005 Japan Science and Tech Corp(JST)  
File 203:AGRIS 1974-2005/Feb  
Dist by NAL, Intl Copr. All rights reserved  
File 342:Derwent Patents Citation Indx 1978-05/200537  
(c) 2005 Thomson Derwent

Set Items Description

--- -----

Cost is in DialUnits

? ds

Terminal set to DLINK

? t s6/9/2 3 5 6

Set Items Description

S1	582	VACCIN?/TI AND (FAIL? OR INEFFECT? OR DISAPPOINT? OR INCON- CLUS? OR TOXIC?)/TI AND (CELLS OR CELL OR WHOLE OR VIABLE? OR VIRUS?)/TI
S2	353	RD (unique items)
S3	9	S2 AND (SUPPOSITORY? OR UROGENTIAL? OR GENITOURIN? OR URIN- ARY? OR VAGINA? OR RECTAL? OR ENTRAL?)
S4	157	S2 AND FAIL?
S5	4	S4 AND WHOLE?
S6	11	S2 AND INEFFECT? NOT S5

## WEST Search History

DATE: Wednesday, June 01, 2005

Hide?	<u>Set</u> <u>Name</u>	<u>Query</u>	<u>Hit</u> <u>Count</u>
		<i>DB=USPT; PLUR=YES; OP=AND</i>	
<input type="checkbox"/>	L1	6099853.pn.	1
		<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=AND</i>	
<input type="checkbox"/>	L2	polysorbate.ti,ab,clm. or poly-sorbate.ti,ab,clm. or \$sorbate.ti,ab,clm.	5837
<input type="checkbox"/>	L3	(peg or pegalate or pegylate or pegylation or pagylating or pegged or pegconjugate or peg\$ or polyethyleneglycol or ethyleneglycol or ethylene-glycol or poly-ethylene-glycol or polyol or polyene or (polymer near5 glycol)).ti,ab,clm.	126905
<input type="checkbox"/>	L4	L3 and l2	530
<input type="checkbox"/>	L5	L3 same l2	325
<input type="checkbox"/>	L6	L5 and (supposit\$ or composition or vaccine or insertion or application or applying)	288
<input type="checkbox"/>	L7	L5 and (supposit\$ or composition or vaccine or insertion or application or applying).ti,ab,clm.	239
<input type="checkbox"/>	L8	L5 same (supposit\$ or composition or vaccine or insertion or application or applying).ti,ab,clm.	160

END OF SEARCH HISTORY

1. 20050096365. 03 Nov 03. 05 May 05. Pharmaceutical compositions with synchronized solubilizer release. Fikstad, David, et al. 514/369; 514/423 514/458 514/469 514/58 A61K031/724 A61K031/426 A61K031/355.

---

☐ 2. 20050096296. 23 Jan 04. 05 May 05. Pharmaceutical compositions with synchronized solubilizer release. Fikstad, David, et al. 514/58; 514/369 514/423 514/458 514/460 514/548 514/571 A61K031/724 A61K031/426 A61K031/401 A61K031/366.

---

☐ 3. 20040209807. 30 Jan 04. 21 Oct 04. Compositions and methods for enhanced mucosal delivery of Y2 receptor-binding peptides and methods for treating and preventing obesity. Quay, Steven C., et al. 514/12; 514/23 514/53 514/58 514/738 A61K038/22 A61K031/70 A61K031/7012 A61K031/724 A61K031/045.

---

☐ 4. 20030138491. 13 Aug 02. 24 Jul 03. Microencapsulation and sustained release of biologically active acid-stable or free sulfhydryl-containing proteins. Tracy, Mark A., et al. 424/486; 424/85.6 A61K038/21 A61K009/14.

---

☐ 5. 20020032171. 08 Jun 01. 14 Mar 02. Clear oil-containing pharmaceutical compositions containing a therapeutic agent. Chen, Feng-Jing, et al. 514/54; 424/727 424/731 424/750 424/757 A61K031/715 A61K035/78.

---

☐ 6. 20020012680. 02 Jul 01. 31 Jan 02. Compositions and methods for improved delivery of lipid regulating agents. Patel, Mahesh V., et al. 424/400; A61K009/00.

---

☐ 7. 6828308. 22 Feb 02; 07 Dec 04. Compositions and methods for the treatment or prevention of inflammation. Mastradonato; Marco, et al. 514/54; 424/486 424/70.15 424/78.36 536/119 536/123.1. A61K031/728 A61K031/715 A61K031/79.

---

☐ 8. 6761903. 08 Jun 01; 13 Jul 04. Clear oil-containing pharmaceutical compositions containing a therapeutic agent. Chen; Feng-Jing, et al. 424/451; 424/43 424/433 424/436 424/441 424/445 424/455 424/456 424/458 424/463 424/464 424/465 424/489 424/490 424/725 514/772.2 514/772.3 514/777 514/779 514/781 514/783 514/784 514/785 514/786 514/937 514/944. A61K009/08 A61K009/10 A61K009/14 A61K009/20 A61K009/48.

---

☐ 9. 6565888. 23 Aug 00; 20 May 03. Methods and compositions for the targeted delivery of biologically active agents. Tracy; Mark A., et al. 424/489; 424/130.1 424/184.1 424/204.1 424/234.1 424/237.1 424/244.1 424/422 424/426 424/490 424/491 424/493 424/497 424/499. A61K009/14.

---

☐ 10. 6465425. 10 Feb 00; 15 Oct 02. Microencapsulation and sustained release of biologically active acid-stable or free sulfhydryl-containing proteins. Tracy; Mark A., et al. 514/2; 514/1 514/724. A01N037/18.

---

☐ 11. 6383471. 06 Apr 99; 07 May 02. Compositions and methods for improved delivery of ionizable hydrophobic therapeutic agents. Chen; Feng-Jing, et al. 424/45; 424/401 424/436 424/451 424/46 514/944. A61K009/12.

---

☐ 12. 6267985. 30 Jun 99; 31 Jul 01. Clear oil-containing pharmaceutical compositions. Chen; Feng-Jing, et al. 424/451; 424/43 424/433 424/436 424/441 424/443 424/455 424/456 424/458 424/463 424/464 424/465 424/489 424/490 424/731 424/735 424/750 424/757 424/764 514/44 514/772.2 514/772.3 514/777 514/778 514/779 514/781 514/783 514/784 514/785 514/786 514/937 514/944. A61K009/08 A61K009/10 A61K009/14 A61K009/20 A61K009/48.

13. JP361225121A. 29 Mar 85. 06 Oct 86. PREVENTIVE FOR ASTHMA AND PREPARATION THEREOF. ABE, TORU, et al. A61K031/41; A61K009/00 A61K009/14 A61K009/16 A61K009/20 A61K009/48 A61K009/70 A61K047/00 C07D257/04.

"PATNO\_JP356150015A" 14. JP 56150015A. Antiallergic agents - esp. effective in treating slow reacting substances of anaphylaxis. A61K031/12 C07C069/95.

[Generate Collection](#)[Print](#)

Terms	Documents
L1 and suppos\$	14

[Prev Page](#)   [Next Page](#)   [Go to Doc#](#)

PUB-NO: JP361225121A

DOCUMENT-IDENTIFIER: JP 61225121 A

TITLE: PREVENTIVE FOR ASTHMA AND PREPARATION THEREOF

PUBN-DATE: October 6, 1986

## INVENTOR-INFORMATION:

NAME	COUNTRY
ABE, TORU	
YANAGIHARA, YUKIYOSHI	
SHINODA, TAKAO	
KONO, SHIGEKATSU	
OHATA, KATSUYA	
OGASAWARA, YOSHIAKI	
KAGEYAMA, SHINJI	
OGUMA, TORU	
TSURITANI, MASAHIRO	
KURODA, TOSHIO	
HASHIMOTO, MITSUMASA	

## ASSIGNEE-INFORMATION:

NAME	COUNTRY
WAKAMOTO PHARMACEUT CO LTD	

APPL-NO: JP60063320

APPL-DATE: March 29, 1985

INT-CL (IPC): A61K 31/41; A61K 9/00; A61K 9/14; A61K 9/16; A61K 9/20; A61K 9/48; A61K 9/70; A61K 47/00; C07D 257/04

## ABSTRACT:

**PURPOSE:** To provide a preventive for asthma, containing a tetrazole derivative as an active component in an amount sufficient to exhibit the effect to suppress the isolation of SRS-A.

**CONSTITUTION:** 5-(3-n-Butyloxalylaminophenyl)tetrazole (abbreviated as MTB) of formula is used as an active component in an amount to exhibit the effect to suppress the isolation of SRS-A. It is mixed uniformly with one or more components selected from polysorbate 80, polyvinyl pyrrolidone, polyethylene glycol, hydroxypropylcellulose and hydroxypropylmethylcellulose to improve the bioavailability of MTB in the presence of a non-aqueous solvent, and the non-aqueous solvent is removed. As an alternative method, MTB is dissolved in liquid PEG and used as an active component of the titled agent optionally after diluting with an inert carrier. It is administered in the form of oral drug such as tablet, granule, etc., or inhalant, suppository, poultice, etc. It is administered at a dose of 10 ~ 500mg per head.

COPYRIGHT: (C)1986,JPO&amp;Japio

## ⑫ 公開特許公報(A)

昭61-225121

⑤ Int. Cl.<sup>4</sup>A 61 K 31/41  
9/00  
9/14

識別記号

A C D

庁内整理番号

④ 公開 昭和61年(1986)10月6日

※審査請求 未請求 発明の数 2 (全8頁)

⑭ 発明の名称 喘息予防剤及びその製剤法

⑰ 特 願 昭60-63320

⑱ 出 願 昭60(1985)3月29日

特許法第30条第1項適用 昭和59年9月30日 日本アレルギー学会事務所発行の「アレルギー第34回  
日本アレルギー学会総会号VOL. 33 NO 9」に発表

⑲ 発 明 者	阿 部	亨	相模原市上鶴間3590 サンキビル506号
⑲ 発 明 者	柳 原	行 義	町田市鶴川5-13-10
⑲ 発 明 者	信 太	隆 夫	相模原市桜台18-1
⑲ 発 明 者	河 野	茂 勝	大津市一里山2-18-21 一里山コーポ105号
⑲ 発 明 者	大 幡	勝 也	宇治市木幡御蔵山39-621
⑲ 発 明 者	小 笠 原	由 明	小田原市栢山1918番地
⑲ 発 明 者	景 山	真 二	神奈川県足柄上郡大井町金手280番地
⑳ 出 願 人	わかもと製薬株式会社		東京都中央区日本橋室町1丁目8番地

最終頁に続く

## 明 細 書

## 1. 発明の名称

喘息予防剤及びその製剤法

## 2. 特許請求の範囲

- (1) SRS-A遊離抑制効果を奏する量の5-(3- $\alpha$ -ブチルオキサリルアミノフェニル)テトラゾールを有効成分として含有することを特徴とする喘息予防剤。
- (2) SRS-A遊離抑制効果を奏する量の5-(3- $\alpha$ -ブチルオキサリルアミノフェニル)テトラゾールをポリソルベート80, ポリビニルピロリドン, ポリオキシエチレン硬化ヒマシ油, ポリエチレングリコール, ヒドロキシメチルセルロース, ヒドロキシプロピルセルロース及びヒドロキシプロピルメチルセルロースの群から選ばれる分散剤の1種又は2種以上と非水溶媒の共存下均一に混合した後非水溶媒を除去するか又は液状ポリエチレング

リコールに溶解した状態で有効成分として含有させることを特徴とする喘息予防剤の製剤法。

## 3. 発明の詳細を説明

## (1) 発明の目的

(産業上の利用分野)

本発明は、喘息予防剤及びその製剤法に関するものである。更に詳しくは、本発明はSRS-A遊離抑制効果を奏する量の5-(3- $\alpha$ -ブチルオキサリルアミノフェニル)テトラゾール(以下MTBという)を有効成分として含有することを特徴とする喘息予防剤を提供するものである。MTBは非常に優れたSRS-A遊離抑制作用とヒスタミン遊離抑制作用を兼備していることから、これを喘息患者に投与すれば、喘息惹起の原因物質である上記両成分の遊離が効果的に抑制され、喘息予防剤としての優れた効果を奏するものである。

(従来の技術)

喘息は発作性の呼吸困難と喘鳴を示す疾患で、気道狭窄によって起こるものである。

気道狭窄の原因としては、気道平滑筋の収縮、気道粘膜の浮腫、気道分枝の亢進、気道内の粘液栓の形成などが挙げられるが、最も重要なのは、気道平滑筋の収縮である。


喘息患者は一般的に気道が過敏であり、吸入アレルギーを含めて多くの抗原に対してIgE抗体を産生し易く、体内にIgE抗体を多量保有しているため、花粉その他のアレルギー(抗原)を吸入した場合、気道粘膜下組織に豊富に存在する肥満細胞の表面で抗原抗体反応が起こり易く、その反応を引金として遊離されるヒスタミン及びSRSAによって、平滑筋収縮を始めとする喘息諸症状を引き起こすものである。〔側ライフ・サイエンス発行；プログレス・イン・メディシン(Progress in Medicine)第3巻第655-666頁(1983年).〕

ヒスタミンによる気管支平滑筋の収縮は非

(3)

本体であり、それらの化学構造も明らかにされている。(免疫薬理第2巻第2号207-213頁, 1984年.)

MTBは本出願人会社の研究部に於いて合成され、抗アレルギー作用を示すことが見出された化合物である。(特開昭57-11975号)



MTBがヒスタミンの遊離を抑制し、抗アレルギー作用を有することはすでに公知であるが、SRSAの遊離抑制作用については全く知られていなかった。〔ジャパニーズ・ジャーナル・オブ・ファーマコロジー(Japanese Journal of Pharmacology)第32巻689-697頁, 1982年〕

(発明が解決しようとする問題点)

本発明者等は、抗喘息剤の開発を目的とし、

(5)

常に敏感、即時的、強力であるが、その重篤な発作は通常~~2-3分程度~~の比較的短時間でおさまる。

(アレルギー第7巻93-104頁, 1958年)

これに対し、SRSAによる気管支平滑筋の収縮は、ゆっくり起こるが強力且つ長時間持続し、喘息患者に堪えられない多大の苦痛を与えるため、このSRSAの遊離を効果的に抑制する薬剤の開発が待望されているところである。〔プログレス・イン・メディシン第3巻655-666頁(1983).〕

SRSA(Slow reacting substance of anaphylaxis)はIgE抗体の関与する抗原抗体反応により肥満細胞等より遊離されるが、ヒスタミン(preformed mediatorに属する)と異なり、その反応の刺激で合成されるもの(newly generated mediatorに属する)であり、アラキドン酸から5-リポキシゲナーゼにより開始される一連の反応により生成されるロイコトリエンC<sub>4</sub>, D<sub>4</sub>, E<sub>4</sub>(LTC<sub>4</sub>, LTD<sub>4</sub>, LTE<sub>4</sub>)がその

(4)

喘息症状を悪化させる最も重要な原因物質がSRSAであることに着目し、SRSAの遊離を効果的に抑制する化合物の探索のため種々スクリーニング試験を行ない、MTBがその目的に合致する化合物であることを見出した。

次いで、MTBの医薬品としての製剤改良研究を行ない、MTB原末が水に難溶性の化合物であり、製剤加工の際非常に取扱いにくく、原末をそのまま通常の方法で製剤化した場合は製剤の崩解性及び溶出性が悪く、有効成分の生体内利用率も不十分であることが分かったので、それらの問題点を解決するため種々検討を行なった結果、このたび喘息予防剤及びその製剤法の発明を完成したものである。

(2) 発明の構成

(問題点を解決するための手段)

本発明は「SRSA遊離抑制効果を奏する量のMTBを有効成分として含有すること」を特徴とする喘息予防剤(以下本薬剤という)

(6)

及びその製剤法」に関するものである。

本薬剤は、通常、錠剤、顆粒剤、散剤、カプセル剤等の経口投与剤として使用されるが、吸入剤、坐剤、パップ剤、注射剤等としても使用出来る。

本薬剤は、通常、成人1回当たり約10～500mg、好ましくは50～300mgの割合で、投与することにより優れた喘息予防効果が期待されるものである。

本薬剤を製造するに当り、慣用の製剤処方を採用することも出来るが、生体内利用率 (Bioavailability) を高めるため、MTBをポリソルベート80、ポリビニルピロリドン (以下PVPという)、ポリオキシエチレン硬化ヒマシ油、ポリエチレングリコール (以下PEGという)、ヒドロキシメチルセルロース、ヒドロキシプロピルセルロース及びヒドロキシプロピルメチルセルロースの群から選ばれる成分 (以下特定分散剤ということがある) の1種又は2種以上と非水溶媒の共存下均一に混

(7)

ロース、コーンスターチ、マンニット、軽質無水ケイ酸、カオリン等が使用出来る。

新製剤法によれば、有効成分MTBが特定分散剤と共に非水溶媒に溶解又は分散された後、溶媒蒸発と共に固溶体を形成するか又は共同沈殿を生じあるいは特定分散剤が液状のPEGの場合はこれにMTBが溶解した状態の製剤が提供され、分散性及び溶出性の著しく優れた生体内利用率の高い本薬剤を製造することが出来る。

示差熱走査熱量測定装置 (DSC30型、島津製作所製) による測定で、MTBは結晶性を示す吸収のピークを示したが、特定分散剤と共に非水溶媒共存下混合処理した物はピークを示さなかったことから、新製剤法に於ける非水溶媒処理は単に混合効率を高めるのみでなく、MTBの結晶構造自体に変化を起こすことが認められた。

又、この新製剤法に於いて、担体を使用せず又は制限して使用してMTBを高濃度に含

(9)

合した後非水溶媒を除去するか、又は液状PEGに溶解した状態で有効成分として含有させ、必要に応じ医薬品として許容される不活性担体で希釈して製剤化するのが好ましい。この方法 (以下新製剤法という) に利用する不活性溶媒はMTBと特定分散剤の両成分を同時に溶解又は分散するもので且つ蒸発除去の容易なものであれば特に限定しないが、通常メタノール、エタノール、イソプロピルアルコール、アセトン、ジクロルメタン等を単独又は混合して使用する。

特定分散剤は水及び非水溶媒の両方に分散又は溶解する比較的高分子の成分からなり、その使用量は種類によりかなり相違するが、MTB1重量部に対し、通常0.01～10重量部の範囲で適宜選択出来る。

PEGは平均分子量200～6000程度のものが好適に使用出来る。

不活性担体は医薬品として許容されるものであれば特に限定されないが、通常結晶セル

(8)

むプレミックスを調製し、このプレミックスを担体で希釈して常法により任意の形態の製剤にすることが出来るし、前記特定分散剤を担体と兼用して多量に使用し製剤化することも出来る。

通常は、一般に用いられる担体、結合剤、甘味料その他補助物質を特定分散剤と共に同時に混合して非水溶媒を用いる湿式造粒法で顆粒剤、散剤、粉剤等を得、必要に応じこれらをカプセルに充填してカプセル剤とするか又は打錠して錠剤にするのが簡便である。

この場合、MTB及び特定分散剤が非水溶媒によって完全には溶解されなかった場合でも、MTB製剤の溶出性は予想外に高い値を示すものである。

#### 実施例 1.

MTB50g、結晶セルロース44g、コーンスターチ10g及び低置換ヒドロキシプロピルセルロース10gを混和し、これにPVP6gをイソ



プロピルアルコール30mlに溶解して添加し、均一に混合した後造粒し、40℃、5時間乾燥後整粒して細粒剤を得る。

#### 実施例 2.

PVP 6gの代りにPVP 4gとポリソルベート-80 2gを使用した外は実施例1と同様にして細粒剤を得、これをゼラチン硬カプセルに充填してカプセル剤を得る。

#### 実施例 3.

MTB 100gをアセトン 500mlに溶解し、これにポリソルベート-80 10gを加えて攪拌混合後、この溶液に結晶セルロース 400g、低置換度ヒドロキシプロピルセルロース 50g及びコーンスターチ 440gを添加し、均一に混合した後、50℃で5時間乾燥後整粒して散剤を得る。

#### 実施例 4.

MTB 100g、結晶セルロース 420g、コー

01

で4時間乾燥して粉末化する。これにステアリン酸マグネシウム 4gを加えて均一に混合し、この混合粉末を4号カプセルに137mg宛充填して、1カプセル中MTB 50mgを含む硬カプセル剤を得る。

#### 実施例 7.

MTB 100gを平均分子量 400 のPEG 400gに溶解し、これに軽質無水ケイ酸 200g及びカルボキシメチルセルロースカルシウム 300gを混合して粉末とする。これを42メッシュの篩を通過させて500mg中MTB 50mg含有の散剤を得る。

#### 実施例 8.

MTB 100g、結晶セルロース 80g、コーンスターチ 20g及び低置換度ヒドロキシプロピルセルロース 16gを混合し、これにPVP 16g及びポリオキシエチレン硬化ヒマシ油 4gをエタノール 120mlに溶解して添加し、均一に混練し

03

ンスターチ 400g及び低置換度ヒドロキシプロピルセルロース 30gを混合した後、PVP 50gをイソプロピルアルコール 550mlに溶解して加え、練合して0.7mm径スクリーンを使って押し出し顆粒状とし、50℃で5時間乾燥後整粒して500mg中50mgのMTBを含有する顆粒剤を得る。

#### 実施例 5.

MTB 10gとヒドロキシプロピルメチルセルロース 20gをジクロルメタン・エタノール等量混液 200mlに加え、溶解後軽質無水ケイ酸 20gを添加して均一に混合し、減圧下溶媒留去乾燥して40メッシュの篩を通過させた後、マンニト 50gを添加して均一に混合し散剤を得る。

#### 実施例 6.

MTB 100gをジクロルメタン・エタノール等量混合液 300mlに溶解し、これにポリソルベート-80 3gを加えて溶解し、次いで結晶セルロース 167gを添加して均一に混合後、50℃

02

で後造粒を行い、これを50℃4時間乾燥し、整粒後タルク 4gを加えて均一に混合し、120mg宛4号カプセルに充填して1カプセル中MTB 50mgを含む硬カプセル剤を得る。

#### 実施例 9.

MTB 100g及び平均分子量 6000 のPEG 80gをジクロルメタン・エタノール等量混合液 500mlに溶解し、これに軽質無水ケイ酸 20gを加え混合後50℃で4時間乾燥後粉碎し、次いで結晶セルロース 88g、カルボキシメチルセルロース 10g及びステアリン酸マグネシウム 2gを加え、常法により打錠して1錠中MTB 約50mg含有する錠剤を得る。

#### 実施例 10.

MTB 100g、結晶セルロース 80g、コーンスターチ 36g及び低置換度ヒドロキシプロピルセルロース 10gを混合し、これにPVP 10g及びポリソルベート-80 2gをエタノール 120

04

mlに溶解して添加し、均一に練合後、0.7mm径スクリーンで押し出し顆粒状とし、50℃で5時間乾燥後整粒し、ステアリン酸マグネシウム2%を加えた後、常法により打錠して直径7mm、重量120mgの錠剤を得る。

#### 実施例 11.

M.T.B 2%をアセトン10%に溶解し、これにポリソルベート-80 0.2%を加えて攪拌混合し、この溶液をカオリン511.8%と均一に混和後乾燥して乾燥粉末を得る。一方、グリセリン440%を100℃に加熱して水分を除き、これにホウ酸45%を加えて溶解し、これに上記乾燥粉末を混和し、冷後チモール0.5%を0.5%のハッカ油に溶解して添加し、よく混和してパップ剤を得る。

#### (3) 発明の効果

本薬剤は、喘息発作の引金となるヒスタミン及びS.R.S-Aの遊離抑制作用が非常に高く、又喘息発作時の気道狭窄（呼吸量低下）

05

(Ro : 対照(未処置)群の遊離量)  
(R : 供試薬剤処置群の遊離量)

供試薬剤は本発明薬剤のM.T.Bと比較薬剤のクロモグリク酸ナトリウム（以下D.S.C.Gという）及びトラニラスト（以下N-5'という）を用い、濃度 $10^{-8}$ ～ $10^{-4}$  (g/ml)の範囲で試験した。なお、ヒスタミンの測定はジャーナル・オブ・アレルギー (Journal of Allergy) 第46巻12～20頁、1970年に記載のメイ・シー・ディー等 (May, C.D. et al) の方法に準じ、螢光法により行った。S.R.S-Aの測定は、薬物学実験（高木敬次郎、小澤光共著、南山堂発行）に記載のマグナス (Magnus) 法により行った。

#### 試験結果

本試験の成績は第1表に示す通りである。各測定値は3回の平均値で示した。

07

の予防効果も非常に優れた喘息の予防薬である。

以下、本薬剤の効果を試験例により詳細に説明する。

#### 試験例 1.

（モルモット肺切片からのヒスタミン及びS.R.S-Aの遊離に及ぼすM.T.Bの抑制効果試験）

#### 試験方法

ハートレイ (Hartley) 系雄性モルモットに抗B.S.Aモルモット血清 (0.5 ml/頭) を静脈内投与して受動的に感作した。2日後、放血致死させて得られた肺を切片とした後、所定濃度の供試薬剤で処置し、5分間経過後抗原を作用させて遊離したヒスタミン及びS.R.S-Aの量を測定し、対照群（薬剤処置しなかった群）の遊離量と比較して遊離抑制率(%)を算出した。

$$\text{遊離抑制率} = \frac{R_o - R}{R_o} \times 100 (\%)$$

06

第 1 表

薬剤濃度 (g/ml)	遊離抑制率 (%)					
	ヒスタミン			S.R.S-A		
	MTB	DSCG	N-5'	MTB	DSCG	N-5'
$10^{-4}$	37	2	16	64	13	54
$10^{-5}$	30	0	5	66	9	11
$10^{-6}$	14			49		
$10^{-7}$	6			28		
$10^{-8}$	8			2		
対照	0	0	0	0	0	0

第1表の成績から明らかなように、本発明薬剤M.T.Bは公知のD.S.C.G及びN-5'に比較してヒスタミン及びS.R.S-Aの遊離抑制率が著しく高く、特にS.R.S-A遊離抑制率は $10^{-6}$ ～ $10^{-4}$  (g/ml)の濃度に於いて49～66%と高い値を示した。

08

## 試験例 2.

アカゲザル肺切片からのヒスタミン及び  
S R S - A の遊離に及ぼす M T B の抑制  
効果試験

## 試験方法

アカゲザルの肺を切片とした後、ヒト・アトピー性血清で受動的に感作を行い、以後試験例 1 と同様な方法で試験し、本発明薬剤 M T B のヒスタミン及び S R S - A 遊離抑制効果を公知の薬剤 D S C G と比較した。

## 試験結果

第 2 表に示す通りである。各測定値は 3 回の平均値で示した。

第 2 表

供試薬剤 濃 度 ( $\mu$ /ml)	抑 制 率 (%)			
	ヒスタミン		S R S - A	
	MTB	DSCG	MTB	DSCG
$10^{-4}$	27	0	16	2
$10^{-5}$	2	0	17	0
$10^{-6}$	12		18	
$10^{-7}$	0		0	
対 照	0	0	0	0

09

第 3 表

供試薬剤 濃 度 ( $\mu$ /ml)	遊 離 抑 制 率 (%)					
	ヒスタミン			S R S - A		
	MTB	DSCG	N-5'	MTB	DSCG	N-5'
$10^{-4}$	27	18	31	47	39	37
$10^{-5}$	23	13	16	47	41	32
$10^{-6}$	23			43		
$10^{-7}$	8			31		
$10^{-8}$	0			2		
対 照	0	0		0	0	0

## 試験例 4. (実験的喘息の予防効果試験)

## 試験方法

モルモット抗 D N P - A S 血清で受動的に感作したモルモットを気密性の箱に入れ、無麻酔、無拘束下、ネブライザーを介して  $2.5 \text{ mg/ml}$  の D N P - B S A を 15 秒間吸入させ実験的喘息を惹起させた。モルモット胸部部の変動を差圧トランデュースを介してレコーダーに記録し、

00

## 試験例 3.

ヒト肺切片からのヒスタミン及び S R S  
- A の遊離に及ぼす M T B の抑制効果試  
験

## 試験方法

肺癌手術により切除されたヒト肺の、肉眼的正常部をアカゲザルの場合に準じてヒト・アトピー性血清で受動的に感作し、以後試験例 1 と同様にして、供試薬剤のヒスタミン及び S R S - A の遊離抑制効果を測定した。

## 試験結果

本試験の結果は第 3 表に示す通りである。各測定値は 3 回の平均値で示した。

00

1 回当たりの呼気量 (tidal volume) を経時的に測定し、喘息惹起前の値と比較して呼気量低下率を求めた。

供試薬剤は、M T B と N - 5' を使用し、それぞれ  $250 \text{ mg/Kg}$  の割合で、喘息惹起の 60 分前に経口投与した。

## 試験結果

本試験の結果は、第 4 表に示す通りである。各測定値は 7 回の平均値で示した。

第 4 表

時 間 (分)	呼 気 量 低 下 率 (%)		
	対 照	M T B	N - 5'
0	0	0	0
1	34	17	26
2	48	17	24
3	47	21	24
4	46	23	26
5	42	21	23
6	41	22	25
8	39	21	28
10	34	21	24
15	30	14	27
20	25	19	22
30	29	15	19

00

第4表の成績から明らかなように、本発明薬剤MTB投与群は薬剤を投与しない対照群及び比較薬剤N-5'投与群のいずれと比較しても、1回呼吸量の低下が非常に少なく、特に、喘息惹起直後の急激な呼吸量低下に対する抑制効果が優れていた。

MTBの温血動物に対する急性毒性は第5表に示す通りである。

第 5 表

投与経路 動物		急性毒性 LD <sub>50</sub> (mg/Kg)			
		i. v.	i. p.	s. c.	p. o.
マウス	雄	1130	1120	>4000	>4000
	雌	1225	1160	>4000	>4000
ラット	雄	1110	1430	>4000	>4000
	雌	1160	1430	>4000	>4000
犬	雄	—	—	—	>4000
	雌	—	—	—	>4000

23

## 試験結果

第6表に示す通り本発明薬剤は著しく高い溶出率を示した。

第 6 表

溶出率(%) 時間(分)	硬カプセル剤		錠 剤	
	本発明A	比較A	本発明B	比較B
5	77.6	3.6	51.2	2.1
10	87.8	6.2	60.0	3.7
20	98.0	12.3	84.3	6.8
30	99.8	17.6	93.6	9.5
40	101.4	22.5	97.8	10.9
60	98.7	31.7	98.4	12.0

本試験に使用した比較薬剤は下記の処方で調製した。

## 比較A

MTB結晶粉末50gを乳糖68g及びステアリン酸マグネシウム2gと均一に混合し、120mgずつカプセルに充填し、1カプセル中MTB

供試動物は次の通りである。

マウス：ddy, 体重 20～23g, 1群20匹  
ラット：wistar, 体重 110～130g, 1群10匹  
犬：beagle, 体重約8kg, 1群3匹

次に新製剤法による本薬剤が著るしく高い溶出率及び生体内利用率(血中濃度)を示すことを説明するため、試験例を示す。

## 試験例 5. (溶出試験)

## 試験方法

溶出液として人工胃液(日本薬局方第1液)500mlを用い、液温37±0.5℃に保ち容器の底に供試薬剤(MTB含量50mg)を入れパドルを100rpmで回転し、所定時間ごとに液をサンプリングし、経時的に液中に溶出されたMTB量を分光光度法により測定した。

## 供試薬剤

本発明A …… 実施例6の硬カプセル剤  
本発明B …… 実施例10の錠剤  
比較A …… 常法による硬カプセル剤  
比較B …… 常法による錠剤

24

50mg含有する硬カプセル剤を得る。

## 比較B

MTB結晶粉末50g、乳糖40g、コーンスターチ15g、結晶セルロース10g、デンプンのり3gを混合し、約40mlの水を加えて練合し、直径7mmのスクリーンより押し出し、顆粒状として乾燥後、整粒してステアリン酸マグネシウム2gを加えて打錠し、1錠中MTB50mg含有する錠剤を得る。

## 試験例 6. (血中濃度測定試験)

## 試験方法

24時間絶食させた体重約10kgの雄性ビーグル犬1群5頭に供試薬剤を経口投与した。薬剤投与後所定時間ごとに採血して、MTBの血中濃度を液体クロマトグラフィーにより測定した。

## 供試薬剤

試験例5と同じ。

## 試験結果

本試験の結果は第7表に示す通りである。

25

26

第 7 表

時間(分)	硬カプセル剤		錠 剤	
	本発明A	比 較 A	本発明B	比 較 B
15	0.23	0.04	0.14	ND
30	0.36	0.08	0.28	0.03
45	0.44	0.12	0.39	0.05
60	0.57	0.14	0.56	0.07
90	0.62	0.30	0.64	0.13
120	0.47	0.18	0.56	0.09
180	0.23	0.08	0.31	0.04
240	0.16	0.05	0.22	0.03
360	0.09	0.03	0.10	ND

備考：1) 表中の成績は5頭の平均値で示した。

2) NDは検出できなかったことを示す。

以上の各試験例から明らかなように、本発明によれば、非常に優れたSR5-A遊離抑制効果を示す喘息予防剤を製造し得るものである。

特許出願人

わかもと製薬株式会社

07

第1頁の続き

⑤Int.Cl.<sup>4</sup>

識別記号

庁内整理番号

A 61 K 9/16

9/20

9/48

9/70

47/00

// C 07 D 257/04

6742-4C

6664-4C

⑦発明者 小 熊

徹

秦野市尾尻450番地の62

⑦発明者 釣 谷

昌 敏

秦野市北矢名666-278

⑦発明者 黒 田

敏 男

相模原市相模台5-12-4-201

⑦発明者 橋 本

光 正

川崎市高津区子母口852-4

L2: Entry 4 of 14

File: PGPB

Jul 24, 2003

PGPUB-DOCUMENT-NUMBER: 20030138491  
PGPUB-FILING-TYPE: new  
DOCUMENT-IDENTIFIER: US 20030138491 A1

TITLE: Microencapsulation and sustained release of biologically active acid-stable or free sulfhydryl-containing proteins

PUBLICATION-DATE: July 24, 2003

## INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Tracy, Mark A.	Arlington	MA	US	
Ward, Kevin L.	Arlington	MA	US	
Scher, David S.	Hudson	MA	US	
Johnson, J. Keith	Hudson	MA	US	

## ASSIGNEE-INFORMATION:

NAME	CITY	STATE	COUNTRY	TYPE CODE
Alkermes Controlled Therapeutics, Inc.	Cambridge	MA		02

APPL-NO: 10/ 217953 [PALM]  
DATE FILED: August 13, 2002

## RELATED-US-APPL-DATA:

Application 10/217953 is a continuation-of US application 09/501934, filed February 10, 2000, US  
Patent No. 6465425

INT-CL: [07] A61 K 38/21, A61 K 9/14

US-CL-PUBLISHED: 424/486; 424/85.6

US-CL-CURRENT: 424/486; 424/85.6

REPRESENTATIVE-FIGURES: NONE

## ABSTRACT:

This invention relates to sustained release compositions, and methods of forming and using said compositions, for the sustained release of biologically active acid-stable or free sulfhydryl-containing proteins, in particular .beta.-IFN. The sustained release composition of this invention comprises a biocompatible polymer having dispersed therein a stabilized biologically active acid-stable or free sulfhydryl-containing protein formulation and a nonionic polymer surfactant.

The method of the invention, for forming a composition for the sustained release of biologically active acid-stable or free sulfhydryl-containing protein, in particular .beta.-IFN, includes dissolving a polymer in a polymer solvent to form a polymer solution, adding a stabilized biologically active acid-stable or

free sulfhydryl-containing protein formulation and a nonionic surfactant to the polymer solution, and then solidifying the polymer to form a polymer matrix containing the stabilized biologically active acid-stable or free sulfhydryl-containing protein formulation, and a nonionic surfactant.

The method of using the sustained release composition of the present invention comprises providing a therapeutically effective blood level of biologically active acid-stable or free sulfhydryl-containing protein in a subject for a sustained period by administering to the subject a dose of the sustained release composition described herein.

#### RELATED APPLICATION

[0001] This application is a continuation of U.S. application Ser. No. 09/501,934, filed Feb. 10, 2000. The entire teachings of the above application are incorporated herein by reference.

DOCUMENT-IDENTIFIER: US 20030138491 A1

TITLE: Microencapsulation and sustained release of biologically active acid-stable or free sulfhydryl-containing proteins

Detail Description Paragraph:

[0059] The composition of this invention can be administered to a human, or other animal, by injection, implantation (e.g., subcutaneously, intramuscularly, intraperitoneally, intracranially, and intradermally), administration to mucosal membranes (e.g., intranasally, intravaginally, intrapulmonary or by means of a suppository), or in situ delivery (e.g., by enema or aerosol spray) to provide the desired dosage of biologically active acid-stable or free sulfhydryl-containing protein, for example .beta.-IFN, based on the known parameters for treatment with such protein of the various medical conditions.

CLAIMS:

10. The sustained release composition of claim 1 wherein the surfactant is selected from the group consisting of: poloxamers, polysorbates, polyethyleneglycols, polyoxyethylene fatty acid esters, bile salts, benzalkonium chloride, polyoxyethylene (40) monostearate and combinations thereof.

21. The sustained release composition of claim 20 wherein the nonionic surfactant is selected from the group consisting of: poloxamers, polysorbates, polyethyleneglycol, polyoxyethylene fatty acid esters and combinations thereof.



PGPUB-DOCUMENT-NUMBER: 20020032171  
PGPUB-FILING-TYPE: new  
DOCUMENT-IDENTIFIER: US 20020032171 A1

TITLE: Clear oil-containing pharmaceutical compositions containing a therapeutic agent

PUBLICATION-DATE: March 14, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Chen, Feng-Jing	Salt Lake City	UT	US	
Patel, Mahesh V.	Salt Lake City	UT	US	
Fikstad, David T.	Salt Lake City	UT	US	

APPL-NO: 09/ 877541 [PALM]

DATE FILED: June 8, 2001

RELATED-US-APPL-DATA:

Application 09/877541 is a continuation-in-part-of US application 09/345615, filed June 30, 1999, US Patent No. 6267985

Application 09/877541 is a continuation-in-part-of US application 09/751968, filed December 29, 2000, PENDING

Application 09/751968 is a continuation-in-part-of US application 09/375636, filed August 17, 1999, US Patent No. 6309663

INT-CL: [07] A61 K 31/715, A61 K 35/78

US-CL-PUBLISHED: 514/54; 424/727, 424/731, 424/750, 424/757

US-CL-CURRENT: 514/54; 424/727, 424/731, 424/750, 424/757

ABSTRACT:

The present invention relates to pharmaceutical compositions and methods for improved solubilization of triglycerides and improved delivery of therapeutic agents. Compositions of the present invention include a carrier, where the carrier is formed from a combination of a triglyceride and at least two surfactants, at least one of which is hydrophilic. Upon dilution with an aqueous medium, the carrier forms a clear, aqueous dispersion of the triglyceride and surfactants.

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation-in-part of U.S. patent application Ser. No. 09/345,615, filed Jun. 30, 1999, and a continuation-in-part of U.S. application Ser. No. 09/751,968, filed Dec. 29, 2000, which is a continuation-in-part of U.S. application Ser. No. 09/375,636, filed Aug. 17, 1999, the disclosures of which are hereby incorporated by reference.

DOCUMENT-IDENTIFIER: US 20020032171 A1

TITLE: Clear oil-containing pharmaceutical compositions containing a therapeutic agent

Summary of Invention Paragraph:

[0024] In another embodiment, the present invention relates to dosage forms comprising the pharmaceutical compositions described herein. The dosage forms may be processed by techniques selected from the group consisting of lyophilization, encapsulation, extruding, compression, melting, molding, spraying, coating, comminution, mixing, homogenization, sonication, granulation, and combinations thereof. Dosage forms include, but are not limited, those selected from the group consisting of pills, capsules, caplets, tablets, granules, beads, powders, solutions, suspensions, emulsions, creams, ointments, lotions, suppositories, sprays, aerosols, pastes, gels, drops, douches, ovules, wafers, troches, cachets, syrups and elixirs.

Detail Description Paragraph:

[0193] Although formulations specifically suited to oral administration are presently preferred, the compositions of the present invention can also be formulated for topical, transdermal, buccal, ocular, pulmonary, vaginal, rectal, transmucosal or parenteral administration, as well as for oral administration. Thus, the dosage form can be a solution, suspension, emulsion, cream, ointment, lotion, suppository, spray, aerosol, paste, gel, drops, douche, ovule, wafer, troche, cachet, syrup, elixir, or other dosage form, as desired. If formulated as a suspension, the composition can further be processed in capsule form.

## CLAIMS:

14. The pharmaceutical composition of claim 6, wherein the hydrophilic surfactant is selected from the group consisting of PEG-10 laurate, PEG-12 laurate, PEG-20 laurate, PEG-32 laurate, PEG-32 dilaurate, PEG-12 oleate, PEG-15 oleate, PEG-20 oleate, PEG-20 dioleate, PEG-32 oleate, PEG-200 oleate, PEG-400 oleate, PEG-15 stearate, PEG-32 distearate, PEG-40 stearate, PEG-100 stearate, PEG-20 dilaurate, PEG-25 glyceryl trioleate, PEG-32 dioleate, PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-20 glyceryl stearate, PEG-20 glyceryl oleate, PEG-30 glyceryl oleate, PEG-30 glyceryl laurate, PEG-40 glyceryl laurate, PEG-40 palm kernel oil, PEG-50 hydrogenated castor oil, PEG-40 castor oil, PEG-35 castor oil, PEG-60 castor oil, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PEG-60 corn oil, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polyglyceryl-10 laurate, PEG-30 cholesterol, PEG-25 phyto sterol, PEG-30 soya sterol, PEG-20 trioleate, PEG-40 sorbitan oleate, PEG-80 sorbitan laurate, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, POE-20 oleyl ether, POE-20 stearyl ether, tocopheryl PEG-100 succinate, PEG-24 cholesterol, polyglyceryl-10 oleate, Tween 40, Tween 60, sucrose monostearate, sucrose monolaurate, sucrose monopalmitate, PEG 10-100 nonyl phenol series, PEG 15-100 octyl phenol series, a poloxamer, and mixtures thereof.

15. The pharmaceutical composition of claim 6, wherein the hydrophilic surfactant is selected from the group consisting of PEG-20 laurate, PEG-20 oleate, PEG-35 castor oil, PEG-40 palm kernel oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, polyglyceryl-10 laurate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, PEG-30 cholesterol, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, PEG-24 cholesterol, sucrose monostearate, sucrose monolaurate, a poloxamer, and mixtures thereof.

16. The pharmaceutical composition of claim 6, wherein the hydrophilic surfactant is selected from the group consisting of PEG-35 castor oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polysorbate 20, polysorbate 80, tocopheryl PEG-1000 succinate, PEG-24 cholesterol, a poloxamer, and mixtures thereof.

thereof.

77. A dosage form comprising the pharmaceutical composition of claim 1, wherein the dosage form is selected from the group consisting of a solution, suspension, emulsion, cream, ointment, lotion, suppository, spray, aerosol, paste, gel, drops, douche, ovule, wafer, troche, cachet, syrup and elixir.

101. The pharmaceutical composition of claim 93, wherein the hydrophilic surfactant is selected from the group consisting of PEG-10 laurate, PEG-12 laurate, PEG-20 laurate, PEG-32 laurate, PEG-32 dilaurate, PEG-12 oleate, PEG-15 oleate, PEG-20 oleate, PEG-20 dioleate, PEG-32 oleate, PEG-200 oleate, PEG-400 oleate, PEG-15 stearate, PEG-32 distearate, PEG-40 stearate, PEG-100 stearate, PEG-20 dilaurate, PEG-25 glyceryl trioleate, PEG-32 dioleate, PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-20 glyceryl stearate, PEG-20 glyceryl oleate, PEG-30 glyceryl oleate, PEG-30 glyceryl laurate, PEG-40 glyceryl laurate, PEG-40 palm kernel oil, PEG-50 hydrogenated castor oil, PEG-40 castor oil, PEG-35 castor oil, PEG-60 castor oil, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PEG-60 corn oil, PEG-6 caprate/caprylate glycerides, PEG-8 capratelcaprylate glycerides, polyglyceryl-10 laurate, PEG-30 cholesterol, PEG-25 phyto sterol, PEG-30 soya sterol, PEG-20 trioleate, PEG-40 sorbitan oleate, PEG-80 sorbitan laurate, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, POE-20 oleyl ether, POE-20 stearyl ether, tocopheryl PEG-100 succinate, PEG-24 cholesterol, polyglyceryl-10 oleate, Tween 40, Tween 60, sucrose monostearate, sucrose monolaurate, sucrose monopalmitate, PEG 10-100 nonyl phenol series, PEG 15-100 octyl phenol series, a poloxamer, and mixtures thereof.

102. The pharmaceutical composition of claim 93, wherein the hydrophilic surfactant is selected from the group consisting of PEG-20 laurate, PEG-20 oleate, PEG-35 castor oil, PEG-40 palm kernel oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, polyglyceryl-10 laurate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, PEG-30 cholesterol, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, PEG-24 cholesterol, sucrose monostearate, sucrose monolaurate, a poloxamer, and mixtures thereof.

103. The pharmaceutical composition of claim 93, wherein the hydrophilic surfactant is selected from the group consisting of PEG-35 castor oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polysorbate 20, polysorbate 80, tocopheryl PEG-1000 succinate, PEG-24 cholesterol, a poloxamer, and mixtures thereof.

157. A dosage form comprising the pharmaceutical composition of claim 88, wherein the dosage form is selected from the group consisting of a solution, suspension, emulsion, cream, ointment, lotion, suppository, spray, aerosol, paste, gel, drops, douche, ovule, wafer, troche, cachet, syrup and elixir.

179. The method of claim 164, wherein the dosage form is selected from the group consisting of a solution, suspension, emulsion, cream, ointment, lotion, suppository, spray, aerosol, paste, gel, drops, douche, ovule, wafer, troche, cachet, syrup and elixir.

First Hit   Fwd Refs

L2: Entry 8 of 14

File: USPT

Jul 13, 2004

DOCUMENT-IDENTIFIER: US 6761903 B2

TITLE: Clear oil-containing pharmaceutical compositions containing a therapeutic agent

Brief Summary Text (26):

In another embodiment, the present invention relates to dosage forms comprising the pharmaceutical compositions described herein. The dosage forms may be processed by techniques selected from the group consisting of lyophilization, encapsulation, extruding, compression, melting, molding, spraying, coating, comminution, mixing, homogenization, sonication, granulation, and combinations thereof. Dosage forms include, but are not limited, those selected from the group consisting of pills, capsules, caplets, tablets, granules, beads, powders, solutions, suspensions, emulsions, creams, ointments, lotions, suppositories, sprays, aerosols, pastes, gels, drops, douches, ovules, wafers, troches, cachets, syrups and elixirs.

Detailed Description Text (155):

Although formulations specifically suited to oral administration are presently preferred, the compositions of the present invention can also be formulated for topical, transdermal, buccal, ocular, pulmonary, vaginal, rectal, transmucosal or parenteral administration, as well as for oral administration. Thus, the dosage form can be a solution, suspension, emulsion, cream, ointment, lotion, suppository, spray, aerosol, paste, gel, drops, douche, ovule, wafer, troche, cachet, syrup, elixir, or other dosage form, as desired. If formulated as a suspension, the composition can further be processed in capsule form.

## CLAIMS:

14. The pharmaceutical composition of claim 6, wherein the hydrophilic surfactant is selected from the group consisting of PEG-10 laurate, PEG-12 laurate, PEG-20 laurate, PEG-32 laurate, PEG-32 dilaurate, PEG-12 oleate, PEG-15 oleate, PEG-20 oleate, PEG-20 dioleate, PEG-32 oleate, PEG-200 oleate, PEG-400 oleate, PEG-15 stearate, PEG-32 distearate, PEG-40 stearate, PEG-100 stearate, PEG-20 dilaurate, PEG-25 glyceryl trioleate, PEG-32 dioleate, PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-20 glyceryl stearate, PEG-20 glyceryl oleate, PEG-30 glyceryl oleate, PEG-30 glyceryl laurate, PEG-40 glyceryl laurate, PEG-40 palm kernel oil, PEG-50 hydrogenated castor oil, PEG-40 castor oil, PEG-35 castor oil, PEG-60 castor oil, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PEG-60 corn oil, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polyglyceryl-10 laurate, PEG-30 cholesterol, PEG-25 phyto sterol, PEG-30 soya sterol, PEG-20 trioleate, PEG-40 sorbitan oleate, PEG-80 sorbitan laurate, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, POE-20 oleyl ether, POE-20 stearyl ether, tocopheryl PEG-100 succinate, PEG-24 cholesterol, polyglyceryl-10 oleate, Tween 40, Tween 60, sucrose monostearate, sucrose monolaurate, sucrose monopalmitate, PEG 10-100 nonyl phenol series, PEG 15-100 octyl phenol series, a poloxamer, and mixtures thereof.

15. The pharmaceutical composition of claim 6, wherein the hydrophilic surfactant is selected from the group consisting of PEG-20 laurate, PEG-20 oleate, PEG-35 castor oil, PEG-40 palm kernel oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, polyglyceryl-10 laurate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, PEG-30 cholesterol, polysorbate 20,

polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, PEG-24 cholesterol, sucrose monostearate, sucrose monolaurate, a poloxamer, and mixtures thereof.

16. The pharmaceutical composition of claim 6, wherein the hydrophilic surfactant is selected from the group consisting of PEG-35 castor oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polysorbate 20, polysorbate 80, tocopheryl PEG-1000 succinate, PEG-24 cholesterol, a poloxamer, and mixtures thereof.

77. A dosage form comprising the pharmaceutical composition of claim 1, wherein the dosage form is selected from the group consisting of a solution, suspension, emulsion, cream, ointment, lotion, suppository, spray, aerosol, paste, gel, drops, douche, ovule, wafer, troche, cachet, syrup and elixir.

103. The method of claim 88, wherein the dosage form is selected from the group consisting of a solution, suspension, emulsion, cream, ointment, lotion, suppository, spray, aerosol, paste, gel, drops, douche, ovule, wafer, troche, cachet, syrup and elixir.

US-PAT-NO: 6465425

DOCUMENT-IDENTIFIER: US 6465425 B1

TITLE: Microencapsulation and sustained release of biologically active acid-stable or free sulfhydryl-containing proteins

DATE-ISSUED: October 15, 2002

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Tracy; Mark A.	Arlington	MA		
Ward; Kevin L.	Arlington	MA		
Scher; David S.	Hudson	MA		
Johnson; J. Keith	Hudson	MA		

## ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE CODE
Alkermes Controlled Therapeutics, Inc.	Cambridge	MA			02

APPL-NO: 09/ 501934 [PALM]

DATE FILED: February 10, 2000

INT-CL: [07] A01 N 37/18

US-CL-ISSUED: 514/2; 514/1, 514/724

US-CL-CURRENT: 514/2; 514/1, 514/724

FIELD-OF-SEARCH: 514/1, 514/2, 514/724

PRIOR-ART-DISCLOSED:

## U.S. PATENT DOCUMENTS

Search Selected

Search ALL

Clear

	PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<input type="checkbox"/>	<u>4252791</u>	February 1981	Grossberg et al.	424/85
<input type="checkbox"/>	<u>4389330</u>	June 1983	Tice et al.	427/213.36
<input type="checkbox"/>	<u>4462940</u>	July 1984	Hanisch et al.	260/112R
<input type="checkbox"/>	<u>4711782</u>	December 1987	Okada et al.	424/455
<input type="checkbox"/>	<u>4767628</u>	August 1988	Hutchinson	424/426
<input type="checkbox"/>	<u>4818542</u>	April 1989	DeLuca et al.	424/491
<input type="checkbox"/>	<u>4853218</u>	August 1989	Yim et al.	424/85.7
<input type="checkbox"/>	<u>4871538</u>	October 1989	Yim et al.	424/85.7

<input type="checkbox"/> <u>4962091</u>	October 1990	Eppstein et al.	514/2
<input type="checkbox"/> <u>4992271</u>	February 1991	Fernandes et al.	424/85.2
<input type="checkbox"/> <u>5004605</u>	April 1991	Hershenson et al.	424/85.6
<input type="checkbox"/> <u>5019400</u>	May 1991	Gombotz et al.	424/497
<input type="checkbox"/> <u>5126147</u>	June 1992	Silvestri et al.	424/497
<input type="checkbox"/> <u>5176907</u>	January 1993	Leong	424/78.08
<input type="checkbox"/> <u>5316774</u>	May 1994	Eury et al.	424/501
<input type="checkbox"/> <u>5441734</u>	August 1995	Reichert et al.	424/85.7
<input type="checkbox"/> <u>5594091</u>	January 1997	Igari et al.	528/271
<input type="checkbox"/> <u>5643566</u>	July 1997	Hanisch et al.	424/85.4
<input type="checkbox"/> <u>5656297</u>	August 1997	Bernstein et al.	424/484
<input type="checkbox"/> <u>5711968</u>	January 1998	Tracy et al.	424/487
<input type="checkbox"/> <u>5874479</u>	February 1999	Martin	514/724
<input type="checkbox"/> <u>5922253</u>	July 1999	Herbert et al.	264/5

## FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
0123291	April 1984	EP	
0215658	September 1986	EP	
0270799	October 1987	EP	
0 251 476	January 1988	EP	
0281299	February 1988	EP	
0307097	August 1988	EP	
0330180	February 1989	EP	
2 620 621	March 1989	FR	
WO 90/09166	August 1990	WO	
WO 90/13285	November 1990	WO	
WO 90/13780	November 1990	WO	
WO 91/12882	September 1991	WO	
WO 91/18927	December 1991	WO	
WO 94/19373	September 1994	WO	
WO 96/03116	February 1996	WO	
WO 96/28143	September 1996	WO	
WO 96/40074	December 1996	WO	
WO 97/13502	April 1997	WO	
WO 98/27963	July 1998	WO	
WO 99/24061	May 1999	WO	

## OTHER PUBLICATIONS

Costantino, H.R., et al., "Solid-Phase Aggregation of Proteins under Pharmaceutically Relevant Conditions," *Journal of Pharmaceutical Sciences*, 83 (12) :1662-1669 (1994).

Constantino, H.R., et al., "Aggregation of a Lyophilized Pharmaceutical Protein, Recombinant Human Albumin. Effect of Moisture and Stabilization by Excipients," *Biotechnology*, 13:493-496 (1995).

Liu, W.R., et al., "Moisture-Induced Aggregation of Lyophilized Proteins in the Solid State," *Biotechnology and Bioengineering*, 37:177-184 (1991).

Lu, W., et al., "Protein Release from Poly(lactic-co-glycolic acid) Microspheres: Protein Stability Problems," *PDA Journal of Pharmaceutical Science & Technology*, 49 (1):13-19 (1995).

Cleland, J.L., et al., "In Vitro Release of Bioactive Recombinant Human Interferon-.gamma. From PLGA Microspheres," *Proceed. Intern. Symp. Control. Rel. Bioac. Mater.*, 22:518-519 (1995).

Phillips, L., "Beta-Interferon: An Exciting New Agent for Multiple Sclerosis," *CPJ/RPC*, 237, 240, 242 and 252 (Jun. 1994).

Gekko, K., et al., "Compressibility-Structure Relationship of Globular Proteins," *Biochemistry*, 25:6563-6571 (1986).

Hanley, D.F., et al., "Pharmacology of Interferons I. Pharmacologic Distinctions Between Human Leukocyte and Fibroblast Interferons," *J. Immunopharmac.*, 1:219-226 (1979).

Hosoi, K., et al., "Structural Characterization of Fibroblast Human Interferon-.beta..sub.1," *Journal of Interferon Research*, 8 : 375-384 (1998).

Rubinstein, M., "The Structure of Human Interferons," *Biochimica et Biophysica. Acta.*, 695:5-16 (1982).

Yu-Chang, J.W., and Hanson, M.A., "Parenteral Formulations of Proteins and Peptides: Stability and Stabilizers," *Journal of Parenteral Science and Technology*, 42 (2S):S4-S27 (1988).

Billiau, A., et al., "Human Fibroblast Interferon for Clinical Trials: Pharmacokinetics and Tolerability in Experimental Animals and Humans," *Antimicrobial Agents and Chemotherapy*, 16(1):56-63 (1979).

ART-UNIT: 1656

PRIMARY-EXAMINER: Riley; Jezia

ATTY-AGENT-FIRM: Hamilton, Brook, Smith and Reynolds, P.C.

#### ABSTRACT:

This invention relates to sustained release compositions, and methods of forming and using said compositions, for the sustained release of biologically active acid-stable or free sulfhydryl-containing proteins, in particular .beta.-IFN. The sustained release composition of this invention comprises a biocompatible polymer having dispersed therein a stabilized biologically active acid-stable or free sulfhydryl-containing protein formulation and a nonionic polymer surfactant.

53 Claims, 4 Drawing figures



DERWENT-ACC-NO: 1982-02787E  
DERWENT-WEEK: 198202  
COPYRIGHT 2005 DERWENT INFORMATION LTD

TITLE: Antiallergic agents - esp. effective in treating slow reacting substances of anaphylaxis

PATENT-ASSIGNEE: TAKEDA CHEM IND LTD (TAKE)

PRIORITY-DATA: 1980JP-0053128 (April 21, 1980)

Search Selected

Search ALL

Clear

PATENT-FAMILY:

PUB-NO

☐ JP 56150015 A

☐ JP 88051125 B

PUB-DATE

November 20, 1981

October 13, 1988

LANGUAGE

PAGES

MAIN-IPC

004

000

APPLICATION-DATA:

PUB-NO

APPL-DATE

APPL-NO

DESCRIPTOR

JP 56150015A

April 21, 1980

1980JP-0053128

INT-CL (IPC): A61K 31/12; C07C 69/95

ABSTRACTED-PUB-NO: JP 56150015A

BASIC-ABSTRACT:

Antiallergic agents contg. quinones of formula (I) are new (where n is 1-10; R1 is Me or MeOH, or two of them combined each other form -CH=CH-CH=CH-; R2 is H, lower alkyl or (CH2CH:C(Me)CH2)mH (where m is 1-10)). (I) are effective in prevention or treatment of allergic diseases caused by slow reacting substances of anaphylaxis (SRS-A) (e.g. bronchial asthma, allergic rhinitis, urticaria). (I) may be administered orally or parenterally (injection, external application, inhalation) at a dose of 0.1-10 mg/kg/adult a day. (I) may be formulated into capsules, granules, powders, tablets, troches, pills, ointments, syrups, injections, suppositories, aerosols, or inhalations, together with various components such as excipient (e.g. sugar, lactose, glucose, starch, mannitol, sorbitol, cellulose, talc, cyclodextrin), binder (e.g. cellulose, methylcellulose, polyvinylpyrrolidone, gelatin, gum arabic, polyethylene glycol (PEG), sugar, starch), disintegrator (e.g. starch, CMC, CMC-Ca), lubricant (e.g. talc), preservative (e.g. Na benzoate, NaHSO3), suspending agent (e.g. methylcellulose, Al stearate), dispersing agent (e.g. Polysorbate 80, Emergen 408, Emerson 310), solvent (e.g. water), and base (e.g. cacao butter, PEG, Whitpsol, vaseline).

ABSTRACTED-PUB-NO: JP 56150015A

EQUIVALENT-ABSTRACTS:

DERWENT-CLASS: A96 B05

CPI-CODES: A12-V01; B10-A06; B12-A07; B12-D02; B12-K02; B12-L04;

⑩ 日本国特許庁 (JP)

⑪ 特許出願公開

⑫ 公開特許公報 (A)

昭56-150015

⑬ Int. Cl.<sup>3</sup>  
A 61 K 31/12

識別記号  
A B F

庁内整理番号  
6408-4C

⑭ 公開 昭和56年(1981)11月20日

発明の数 1  
審査請求 未請求

(全 4 頁)

⑮ 抗アレルギー剤

豊中市新千里南町2丁目26番3号

⑯ 特 願 昭55-53128

⑰ 発 明 者 渡辺正純

⑱ 出 願 昭55(1980)4月21日

川西市清和台西2丁目4番地の54

⑲ 発 明 者 牧良孝

京都市西京区大原野上里鳥見町5番地の17

⑳ 出 願 人 武田薬品工業株式会社

大阪市東区道修町2丁目27番地

㉑ 発 明 者 寺尾泰次

㉒ 代 理 人 弁理士 松居祥二


明 細 書

1. 発明の名称

抗アレルギー剤

2. 特許請求の範囲

式



(式中、nは1から10までの整数を、R<sup>1</sup>はメチル基、メトキシ基または2つのR<sup>1</sup>で-CH=CH-CH=CH-基を示し、R<sup>2</sup>は水素原子、低級アルキル基または-(CH<sub>2</sub>-CH=C(CH<sub>3</sub>)-CH<sub>2</sub>)<sub>m</sub>-H基(mは1から10までの整数)を示す)で表わされる化合物を含有する抗アレルギー剤。

3. 発明の詳細な説明


本発明は新規抗アレルギー剤に関する。

一般にアレルギーに基づくと考えられる疾患は即時型と遅延型に分類され、前者は抗原抗体反応の結果生体局所に化学的伝達体が産生、放出され、この物質が気管支筋、肺静脈などの平滑筋を収縮

したり皮膚血管の透過性を亢進するなど生体に障害をひきおこすと考えられている。このような化学的伝達体としてヒスタミンや多価不飽和脂肪酸(特にアラキドン酸)のリボキシゲナーゼによる代謝産物、たとえば slow reacting substance of anaphylaxis (以下SRS-Aと略す)などがあげられるが、SRS-Aによるアレルギー疾患に対し有効な化合物は現在広く検討段階にありその報告例は3-amino-1-(m-trifluoromethyl)-phenyl-2-phrazoline(BW-755Cと略す、FEBS Lett., 110巻、213-215頁、1980年)、5, 8, 11, 14-eicosatetraynoic acid(TYAと略す、Prostaglandins, 14巻、21-38頁、1977年)、taicalein phosphate disodium(BPSと略す、代謝、10巻、730-739頁(1973))など極めて少ない。本発明者らは抗アレルギー剤として、なかでもSRS-Aの産生、放出を抑制する抗アレルギー剤の探索を進めてきた。その結

果、下式(Ⅰ)で表わされる化合物がSRS-Aの産生、放出を極めて強力に抑制し、抗アレルギー剤として有用であることを見出し、これに基づいて本発明を完成した。

すなわち、本発明は式




(式中、nは1から10までの整数を、R<sup>1</sup>はメチル基、メトキシ基または2つのR<sup>1</sup>で-CH=CH-CH=CH-基を示し、R<sup>2</sup>は水素原子、低級アルキル基または(CH<sub>2</sub>-CH=C(CH<sub>3</sub>)-CH<sub>2</sub>)<sub>m</sub>H基(mは1から10までの整数)を示す)で表わされる化合物を含有する抗アレルギー剤である。

上記式(Ⅰ)に関し、R<sup>2</sup>で示される低級アルキル基としては、たとえばC<sub>1</sub>-<sub>4</sub>アルキル基(例、メチル、エチル、プロピル、イソプロピル、ブチル、イソブチル、sec-ブチル、tert-ブチル基)があげられる。

上記化合物(Ⅰ)のなかでもR<sup>1</sup>がメトキシ基、


R<sup>2</sup>が水素原子の化合物は上記薬理作用において優れており、キノン側鎖における炭素数が10~20の化合物が本発明の目的に特に適した化合物である。

上記式(Ⅰ)で示されるキノン類は生体内では式



(式中、R<sup>1</sup>、R<sup>2</sup>およびnは前記と同意義)で表わされるヒドロキノン類と相互変換しており、これらの化合物は生理学的意義において同等である。

化合物(Ⅰ)は、たとえば特願昭54-61915号に開示された方法またはそれらに準じて以下のように製造し得る。




(式中、R<sup>3</sup>は水酸基の保護基を、R<sup>2'</sup>は低級アルキル基を示し、他の記号は前記と同意義)

化合物(Ⅰ)は前述のとおりSRS-Aの産生、放出を極めて強力に抑制し、SRS-Aに起因する種々のアレルギー症(例、気管支喘息、アレルギー性鼻炎、じん麻疹)の治療、予防に有用である。

本発明の抗アレルギー剤は、その有効成分である式(Ⅰ)で示される化合物自体をそのまま投与することもできるが、一般には種々の医薬組成物として投与される。このような医薬組成物の剤形の例としては、例えばカプセル剤、顆粒剤、散剤、錠剤、トローチ剤、丸剤、軟膏剤、シロップ剤、注射剤、坐剤、エロゾル剤、吸入剤等が挙げられる。

また医薬組成物に使用されるものとしては、例えば白糖、乳糖、ブドウ糖、でん粉、マンニト、ソルビット、セルロース、タルク、シクロデキストリン等の賦形剤、セルロース、メチルセルロース、ポリビニルピロリドン、ゼラチン、アラビア




ゴム、ポリエチレングリコール、白糖、でん粉等の結合剤、でん粉、カルボキシメチルセルロース、カルボキシメチルセルロースのカルシウム塩等の崩壊剤、タルク等の滑沢剤、安息香酸ナトリウム、亜硫酸水素ナトリウム等の保存剤、メチルセルロース、ステアリン酸アルミニウム等のけんだく化剤、ポリソルベート80、エマルゲン408、エマゾール310等の分散剤、水等の溶剤、カカオ脂、ポリエチレングリコール、ウエテアソール、白色ワセリン等の基剤等が挙げられ、これらは製剤の種類に応じて適宜選択される。

この発明の抗アレルギー剤は通常の抗アレルギー剤に準じて経口的または非経口的（例、注射、塗布、吸入等）に人間を含む哺乳動物に投与され、皮膚症状、喘息および鼻炎等のアレルギー症状等を効果的に抑制することができる。

本発明の抗アレルギー剤の毒性は極めて低く、臨床的な投与量としては経口もしくは非経口のルートにより、成人1日当たり化合物(Ⅰ)として約0.1-10mg/kg程度が好んで用いられる。

第一表

SRS-Aの産生、放出に対する抑制作用



化合物*)			濃度 (μM)	抑制作用 (%)
R <sup>1</sup>	R <sup>2</sup>	n		
CH <sub>3</sub> O	H	2	1	35.0±6.3
CH <sub>3</sub> O	H	3	1	21.8±8.6
CH <sub>3</sub> O	H	3	1	24.3±14.2
BW-755C			10	4.7±3.7
			100	15.3±4.3
TYA			1	1.3±1.2
			10	56.9±5.9
BPS			10	18.7±6.6
			100	50.8±10.7

\*) 化合物はエタノールまたはジメチルスルホキシド溶液として添加した。

\*\*) 抗原抗体反応によるSRS-Aの産生、放

## 実験例1

本発明の抗アレルギー剤のSRS-A産生に対する抑制作用をOrangeおよびMooreの方法〔J. Immunol., 116巻, 392頁(1976年)〕に従って測定した。すなわち、抗原として卵白アルブミンを用いて感作したモルモット(Hartley 系雄雄, 体重300-350g)の肺切片に、化合物(Ⅰ)と抗原を同時に添加し、その際産生、放出されるSRS-A量をBrocklehurst (J. physiol., 151巻, 416-435頁, 1960年)の方法によつて測定した。その結果第一表に示すように本件化合物は低濃度においてSRS-Aの産生、放出を強力に抑制し、その強度は前記公知化合物BW-755C, TYA, BPSなどに較べて顕著に優れていることが認められた。

出の阻害率。

## 実施例1

カプセル剤

本発明化合物(Ⅰ)式中R<sup>1</sup>=CH<sub>3</sub>O, R<sup>2</sup>=H,

n=3) 30mg

微結晶セルロース 30mg

乳糖 57mg

ステアリン酸マグネシウム 3mg

全量 120mg

上記成分を常法により混合したのちゼラチンカプセルに充填しカプセル剤とした。

## 実施例2

錠剤

本発明化合物(Ⅰ)式中R<sup>1</sup>=CH<sub>3</sub>O, R<sup>2</sup>=H,

n=2) 30mg

乳糖 44mg

でん粉 10.6mg

でん粉(のり用) 5mg

ステアリン酸マグネシウム 0.4mg

カルボキシメチルセルロースカルシウム 20mg

全量 110 ㍑

上記成分を常法により混合したのを錠剤とした。

### 実施例3

軟カプセル剤

本発明化合物(式中  $R = \text{C}_n\text{H}_{2n+1}$ ,  
 $n=3$ ) 30 ㍑

トウモロコシ油 110 ㍑

全量 140 ㍑

上記成分を混合溶液としたのを常法により軟カ  
 プセル剤とした。

代理人 弁理士 松 居 祥 二

First Hit   Fwd Refs

L8: Entry 101 of 160

File: USPT

Oct 19, 1999

DOCUMENT-IDENTIFIER: US 5968490 A

TITLE: Antiperspirant deodorant compositions

## CLAIMS:

14. The composition of claim 2 wherein the nonionic surfactant comprises methyl gluceth-20, methyl gluceth-10, PEG-20 methyl glucose distearate, PEG-20 methyl glucose sesquistearate, PEG-200 castor oil, C.sub.11-15 pareth-20, ceteth-8, ceteth-12, dodoxynol-12, laureth-15, PEG-20 castor oil, polysorbate 20, steareth-20, polyoxyethylene-10 cetyl ether, polyoxyethylene-10 stearyl ether, polyoxyethylene-20 cetyl ether, polyoxyethylene-21 stearyl ether, polyoxyethylene-10 oleyl ether, polyoxyethylene-20 oleyl ether, an ethoxylated nonylphenol having at least 9 ethylene oxide moieties, an ethoxylated octylphenol having at least 9 ethylene oxide moieties, an ethoxylated dodecyl phenol having at least 9 ethylene oxide moieties, an ethoxylated fatty (C.sub.6 -C.sub.22) alcohol having at least 9 ethylene oxide moieties, polyoxyethylene-20 isohexadecyl ether, dimethicone copolyol, polyoxyethylene-23 glycerol laurate, polyoxyethylene-20 glyceryl stearate, PPG-10 methyl glucose ether, PPG-20 methyl glucose ether, a polyoxyethylene-20 sorbitan monoester, polyoxyethylene-80 castor oil, polyoxyethylene-15 tridecyl ether, polyoxyethylene-6 tridecyl ether, and mixtures thereof.

18. The composition of claim 16 wherein the first surfactant is selected from the group consisting of isoceteth-20, ceteth-20, dimethicone copolyol, an ethoxylated nonylphenol having at least 9 ethylene oxide moieties, an ethoxylated dodecylphenol having at least 9 ethylene oxide moieties, an ethoxylated octylphenol having at least 9 ethylene oxide moieties, steareth-10, PEG-20 glyceryl stearate, steareth-20, POE(6)tridecylether, PEG-80 castor oil, steareth-21, polysorbate 20, an ethoxylated fatty (C.sub.6 -C.sub.22) alcohol having at least 9 ethylene oxide moieties, and mixtures thereof.

First Hit   Fwd Refs

L8: Entry 102 of 160

File: USPT

Oct 5, 1999

DOCUMENT-IDENTIFIER: US 5961997 A

TITLE: Antipruritic composition

## CLAIMS:

1. A hypoallergenic, noncomedogenic, nonacnegenic, antipruritic composition which is oil-free, lanolin-free, fragrance-free, and free of Imidazolidinyl urea, quaternium-15, germall, the composition comprising about 0.5% to 1.5% menthol, about 0.5% to 1.5% camphor, about 0.5% to 1.0% phenol, about 1% to 10% lidocaine, about 1% to 2.5% pramoxine, about 10% to 50% distilled water, about 5% to 25% emulsifying wax NF, about 2% to 7% behenric acid, about 1% to 4% PEG 120 methylglucose dioleate, about 1% to 5% polysorbate 20, about 1% to 3% polysorbate 80, about 1% to 5% sorbitan sesquioleate, about 0.5% to 0.15% methyl paraben, about 0.3% to 0.10% propylparaben, about 1% to 5% C12-15 alkyl benzoate in a carrier containing only those ingredients rated 0 or 1 with respect to comedogenicity and irritancy.

First Hit   Fwd Refs

L8: Entry 93 of 160

File: USPT

Aug 6, 2002

DOCUMENT-IDENTIFIER: US 6428816 B1

TITLE: Carotenoid agent for inhibiting the conversion of epithelial cells to tumors

## CLAIMS:

1. A method of inhibiting carcinogen-mediated conversion of melanocyte cells to melanomas, comprising the step of administering to the cells an anti-tumorigenic composition comprising: 0.1 to 10% by weight of a water insoluble carotenoid; and 90 to 99.9% by weight of a non-toxic carrier medium including a suspending agent selected from the group consisting of fatty acids, triglyceride lipids, non-saponifiable lipids, and combinations thereof; an emulsifier selected from the group consisting of a Tween polysorbate glycerol fatty acid esters and acetylated esters of fatty acids; and a water soluble dispersing agent which is a sugar or a polyol.

6. A method of protecting DNA against carcinogen-mediated damage thereby to inhibit conversion of melanocyte cells to melanoma, comprising the step of administering to the cells a composition comprising 0.1 to 10% by weight of a water insoluble carotenoid which includes .mu.-carotene; and 90 to 99.9% by weight of a non-toxic carrier medium including a suspending agent selected from the group consisting of fatty acids, triglyceride lipids, non-saponifiable lipid preparations, soluble hydrocarbons and combinations thereof; an emulsifier selected from the group consisting of a Tween polysorbate glycerol fatty acid esters and acetylated esters of fatty acids; and a water soluble dispersing agent which is a sugar or a polyol.



US-PAT-NO: 6428816

DOCUMENT-IDENTIFIER: US 6428816 B1

TITLE: Carotenoid agent for inhibiting the conversion of epithelial cells to tumors

DATE-ISSUED: August 6, 2002

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Schlupalius; Lance Elliot	Ashwood			AU
Buckmeier; Julie A.	Long Beach	CA		
Meyskens, Jr.; Frank L.	Irvine	CA		

## ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE CODE
Cognis Australia Pty., Ltd.	Broadmeadows			AU	03

APPL-NO: 08/ 718522 [PALM]

DATE FILED: January 16, 1997

## PARENT-CASE:

This application is a continuation-in part of U.S. application Ser. No. 08/743,174, filed Nov. 5, 1996, now U.S. Pat. No. 6,132,790, which is a continuation of U.S. application Ser. No. 08/604,359, filed Feb. 21, 1996, now abandoned, which is a continuation of U.S. application Ser. No. 08/204,188, filed on Apr. 29, 1994, now abandoned, which is a 371 of PCT/AU92/00470 filed on Sep. 7, 1992. Benefit of these applications is claimed pursuant to 35 U.S.C. .sectn.120.

## FOREIGN-APPL-PRIORITY-DATA:

COUNTRY	APPL-NO	APPL-DATE
AU	PM4931	October 8, 1994

## PCT-DATA:

APPL-NO	DATE-FILED	PUB-NO	PUB-DATE	371-DATE	102(E)-DATE
PCT/AU95/00199	April 10, 1995	WO95/27483	Oct 19, 1995	Jan 16, 1997	Jan 16, 1997

INT-CL: [07] A61 K 35/78, A61 K 31/07

US-CL-ISSUED: 424/725; 424/773, 514/725, 514/938

US-CL-CURRENT: 424/725; 424/773, 514/725, 514/938

FIELD-OF-SEARCH: 424/195.1, 424/725, 424/773, 514/725, 514/938

PRIOR-ART-DISCLOSED:

## U.S. PATENT DOCUMENTS

Search Selected

Search ALL

Clear

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<input type="checkbox"/> <u>5230836</u>	July 1993	Todd, Jr.	252/407
<input type="checkbox"/> <u>5705180</u>	January 1998	Schlipalius	424/423
<input type="checkbox"/> <u>6132790</u>	October 2000	Schlipalius	426/540

## FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
93/04598	March 1993	WO	
94/21231	September 1994	WO	

## OTHER PUBLICATIONS

Mathews-Roth, Oncology, 39:33-37, 1982.\*  
 Mathews-Roth, Photochem. and Photobiol., 37, 509-511, 1983.\*  
 Mathews-Roth, Fed. Proc., 46:1890-1893, 1987.\*  
 Ghosh et al., Molecular and Cellular Biology, 13:6992-6999, 1993.\*  
 Georgiadis et al., J. Investigative Dermatology, 98:657, Abstract #631, 1992.\*  
 Rousseau et al., Free Radical Biology and Medicine, 13:407-433, 1992.\*  
 Bain et al., Ann. Epidemiol., 3:235-238, 1993.\*  
 Epstein, Photochem. Photobiol., 25:211-213, 1977.\*  
 van Poppel, Eur. J. Cancer, 29A:1335-1344, 1993.\*  
 "Polysorbates", 7742, Editor Susan budavari et al., The Merck Index An Encyclopedia of Chemicals, Drugs, and Biologicals, 1996, 12th Edition, Merck Research Laboratories Division of Merck & Co., Inc. Whitehouse Station, NJ.  
 Moon RC, "Comparative Aspects of Carotenoids and Retinoids as Chemopreventive Agents for Cancer," J. Nutr. (1989) 119:127-134.  
 Bertram, JS Pung A., Churley M. et al., "Diverse Carotenoids Protect Against Chemically Induced Neoplastic Transformation," Carcinogenesis (1991) 12:671-678.  
 Hazuka MB, Prasad-Edwards J., Newman F., et al., "Beta-carotene Induces Morphological Differentiation and Decreases Adenylate Cyclase Activity in Melanocyte Cells in Culture," J. Am. Coll. Nutr. (1990) 9:143-149.  
 Schultz TD, Chew BP, Seatman WR et al, "Inhibitory Effect of Conjugated Dienoic Derivatives of Linoleic Acid and Beta-carotene on the in vitro Growth of Human Cancer Cells," Canc Letters (1992) 63:125-133.  
 Schwartz JL, Shklar G, "The Selective Cytotoxic Effect of Carotenoids and a-tocopherol on Human Cancer Cell Lines in vitro," J. Oral Maxillofac Surg 0:367-373.  
 Schwartz JL, Tanaka J., Khandekar V, et al, "Beta-Carotene and/or Vitamin E as modulators of Alkylating Agents in SCC-25 Human Squamous Carcinoma Cells," Canc Chemother-Pharmacol (1992) 29:207-213.  
 Zhang L-X, Cooney RV, Bertram JS, "Carotenoids Enhance Gap Junctional Communication and Inhibit Lipid Peroxidation in C3H/10T1/2 Cells; Relationship to Their Cancer Chemopreventive Action," Carcinogenesis (1991) 12:2109-2114.  
 Zhang L-X, Cooney RV, Bertram JS, "Carotenoids Up-regulate Connexin 43 Gene Expression Independent of Their Provitamin A or Antioxidant Properties," Canc Res (1992) 52:5707-5712.  
 Gilchrest BA, Soler NA, Staff JS and Mihm MC Jr., "The Human Sunburn Reaction: Histologic and

Biochemical Studies," J. Am. Acad. Dermatol. (1981) 5:411-422.

Simon, MM et al, "UVB Light Induces Nuclear Factor Kappa B (NF Kappa B) Activity Independently From Chromosomal DNA Damage in Cell-Free Cytosolic Extracts," The Journal of Investigative Dermatology, 102 (1994) 3:422.

ART-UNIT: 1651

PRIMARY-EXAMINER: Tate; Christopher R.

ATTY-AGENT-FIRM: Drach; John E. Ettelman; Aaron R.

**ABSTRACT:**

A method of inhibiting carcinogen mediated conversion of melanocyte cells and protecting DNA against carcinogen-mediated damage by administering a composition comprising a water insoluble carotenoid such as beta-carotene is disclosed.

19 Claims, 18 Drawing figures

DOCUMENT-IDENTIFIER: US 20050020576 A1

TITLE: Kappa agonist compounds and pharmaceutical formulations thereof

Detail Description Paragraph:

[0665] The compositions are formulated as injectables, as oral and rectal formulations for systemic administration, and for local and topical administration as creams, aqueous or non-aqueous suspension, lotions, emulsions, suspensions or emulsions containing micronized particles, gels, foams aerosols, solids and other suitable vehicles for application to the skin, eyes, lips and mucosa, as suppositories or cream for vaginal administration, and as combinations with bandages, patches, bioadhesives and dressings. The compounds may be formulated in combination with other agents, such as local anesthetics and other therapeutic agents. The other agents may be mixed in the compositions are provided and administered prior to, simultaneously with or subsequent to administration of the compositions provided for the methods herein. Such agents include, but are not limited to: antibiotics, including cephalosporins, .beta.-lactams, tetracyclines, vancomycins, sulfas and aminoglycosides; antivirals, including acyclovir; and antifungals including clotrimazole.

Detail Description Paragraph:

[0718] Examples of aqueous vehicles include Sodium Chloride Injection, Ringers Injection, Isotonic Dextrose Injection, Sterile Water Injection, Dextrose and Lactated Ringers Injection. Nonaqueous parenteral vehicles include fixed oils of vegetable origin, cottonseed oil, corn oil, sesame oil and peanut oil. Antimicrobial agents in bacteriostatic or fungistatic concentrations must be added to parenteral preparations packaged in multiple-dose containers which include phenols or cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl p-hydroxybenzoic acid esters, thimerosal, benzalkonium chloride and benzethonium chloride. Isotonic agents include sodium chloride and dextrose. Buffers include phosphate and citrate. Antioxidants include sodium bisulfate. Local anesthetics include procaine hydrochloride. Suspending and dispersing agents include sodium carboxymethylcellulose, hydroxypropyl methylcellulose and polyvinylpyrrolidone. Emulsifying agents include Polysorbate 80 (Tween 80). A sequestering or chelating agent of metal ions include EDTA. Pharmaceutical carriers also include ethyl alcohol, polyethylene glycol and propylene glycol for water miscible vehicles and sodium hydroxide, hydrochloric acid, citric acid or lactic acid for pH adjustment.

DOCUMENT-IDENTIFIER: US 20020012680 A1

TITLE: Compositions and methods for improved delivery of lipid regulating agents

Detail Description Paragraph:

[0137] Although formulations specifically suited to oral administration are presently preferred, the compositions of the present invention can also be formulated for topical, transdermal, ocular, pulmonary, vaginal, rectal, transmucosal or parenteral administration, in the form of a triglyceride-free cream, lotion, ointment, suppository, gel or the like. If such a formulation is desired, other additives may be included, such as are well-known in the art, to impart the desired consistency and other properties to the formulation. The compositions of the present invention can also be formulated as a spray or an aerosol. In particular, the compositions may be formulated as a sprayable solution, and such formulation is particularly useful for spraying to coat a multiparticulate carrier, such as a bead. Such multiparticulate carriers are well known in the art.

## CLAIMS:

13. The pharmaceutical composition of claim 5, wherein the hydrophilic surfactant is PEG-10 laurate, PEG-12 laurate, PEG-20 laurate, PEG-32 laurate, PEG-32 dilaurate, PEG-12 oleate, PEG-15 oleate, PEG-20 oleate, PEG-20 dioleate, PEG-32 oleate, PEG-200 oleate, PEG400 oleate, PEG-15 stearate, PEG-32 distearate, PEG-40 stearate, PEG-100 stearate, PEG-20 dilaurate, PEG-25 glyceryl trioleate, PEG-32 dioleate, PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-20 glyceryl stearate, PEG-20 glyceryl oleate, PEG-30 glyceryl oleate, PEG-30 glyceryl laurate, PEG-40 glyceryl laurate, PEG-40 palm kernel oil, PEG-50 hydrogenated castor oil, PEG-40 castor oil, PEG-35 castor oil, PEG-60 castor oil, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PEG-60 corn oil, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polyglyceryl-10 laurate, PEG-30 cholesterol, PEG-25 phyto sterol, PEG-30 soya sterol, PEG-20 trioleate, PEG-40 sorbitan oleate, PEG-80 sorbitan laurate, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, POE-20 oleyl ether, POE-20 stearyl ether, tocopheryl PEG-100 succinate, PEG-24 cholesterol, polyglyceryl-10 oleate, Tween 40, Tween 60, sucrose monostearate, sucrose monolaurate, sucrose monopalmitate, PEG 10-100 nonyl phenol series, PEG 15-100 octyl phenol series, a poloxamer, or a mixture thereof.

14. The pharmaceutical composition of claim 5, wherein the hydrophilic surfactant is PEG-20 laurate, PEG-20 oleate, PEG-35 castor oil, PEG-40 palm kernel oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, polyglyceryl-10 laurate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, PEG-30 cholesterol, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, PEG-24 cholesterol, sucrose monostearate, sucrose monolaurate, a poloxamer, or a mixture thereof.

15. The pharmaceutical composition of claim 5, wherein the hydrophilic surfactant is PEG-35 castor oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polysorbate 20, polysorbate 80, tocopheryl PEG-1000 succinate, PEG-24 cholesterol, a poloxamer, or a mixture thereof.


67. A dosage form comprising the pharmaceutical composition of claim 1 formulated as a solution, a cream, a lotion, an ointment, a suppository, a spray, an aerosol, a paste or a gel.

83. The pharmaceutical composition of claim 75, wherein the hydrophilic surfactant is PEG-10 laurate, PEG-12 laurate, PEG-20 laurate, PEG-32 laurate, PEG-32 dilaurate, PEG-12 oleate, PEG-15 oleate, PEG-20 oleate, PEG-20 dioleate, PEG-32 oleate, PEG-200 oleate, PEG-400 oleate, PEG-15 stearate,

PEG-32 distearate, PEG-40 stearate, PEG-100 stearate, PEG-20 dilaurate, PEG-25 glyceryl trioleate, PEG-32 dioleate, PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-20 glyceryl stearate, PEG-20 glyceryl oleate, PEG-30 glyceryl oleate, PEG-30 glyceryl laurate, PEG-40 glyceryl laurate, PEG-40 palm kernel oil, PEG-50 hydrogenated castor oil, PEG-40 castor oil, PEG-35 castor oil, PEG-60 castor oil, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PEG-60 corn oil, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polyglyceryl-10 laurate, PEG-30 cholesterol, PEG-25 phyto sterol, PEG-30 soya sterol, PEG-20 trioleate, PEG-40 sorbitan oleate, PEG-80 sorbitan laurate, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, POE-20 oleyl ether, POE-20 stearyl ether, tocopheryl PEG-100 succinate, PEG-24 cholesterol, polyglyceryl-10 oleate, Tween 40, Tween 60, sucrose monostearate, sucrose monolaurate, sucrose monopalmitate, PEG 10-100 nonyl phenol series, PEG 15-100 octyl phenol series, a poloxamer, or a mixture thereof.

84. The pharmaceutical composition of claim 75, wherein the hydrophilic surfactant is PEG-20 laurate, PEG-20 oleate, PEG-35 castor oil, PEG-40 palm kernel oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, polyglyceryl-10 laurate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, PEG-30 cholesterol, polysorbate 20, polysorbate 80, POE-9 lauryl ether, POE-23 lauryl ether, POE-10 oleyl ether, PEG-24 cholesterol, sucrose monostearate, sucrose monolaurate, a poloxamer, or a mixture thereof.

85. The pharmaceutical composition of claim 75, wherein the hydrophilic surfactant is PEG-35 castor oil, PEG-40 hydrogenated castor oil, PEG-60 corn oil, PEG-25 glyceryl trioleate, PEG-6 caprate/caprylate glycerides, PEG-8 caprate/caprylate glycerides, polysorbate 20, polysorbate 80, tocopheryl PEG-1000 succinate, PEG-24 cholesterol, a poloxamer, or a mixture thereof.

 137. The method of claim 136, wherein the dosage form is a capsule, a cream, a lotion, an ointment, a suppository, a paste or a gel.



US 20020012680A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2002/0012680 A1**  
**Patel et al.** (43) **Pub. Date: Jan. 31, 2002**(54) **COMPOSITIONS AND METHODS FOR  
IMPROVED DELIVERY OF LIPID  
REGULATING AGENTS**(76) **Inventors: Mahesh V. Patel, Salt Lake City, UT  
(US); Feng-Jing Chen, Salt Lake City,  
UT (US)**

Correspondence Address:  
**REED & ASSOCIATES**  
**800 MENLO AVENUE**  
**SUITE 210**  
**MENLO PARK, CA 94025 (US)**

(21) **Appl. No.: 09/898,553**(22) **Filed: Jul. 2, 2001****Related U.S. Application Data**(63) Continuation of application No. 09/258,654, filed on  
Feb. 26, 1999, now Pat. No. 6,294,192.**Publication Classification**(51) **Int. Cl.<sup>7</sup> ..... A61K 9/00**  
(52) **U.S. Cl. .... 424/400**(57) **ABSTRACT**

The present invention relates to triglyceride-free pharmaceutical compositions for delivery of hydrophobic therapeutic agents. Compositions of the present invention include a hydrophobic therapeutic agent and a carrier, where the carrier is formed from a combination of a hydrophilic surfactant and a hydrophobic surfactant. Upon dilution with an aqueous solvent, the composition forms a clear, aqueous dispersion of the surfactants containing the therapeutic agent. The invention also provides methods of treatment with hydrophobic therapeutic agents using these compositions.

102(2)?

**INVENTOR-INFORMATION:**

NAME	CITY	STATE	COUNTRY	RULE-47
Gao, Ping	Portage	MI	US	
Morozowich, Walter	Kalamazoo	MI	US	\

APPL-NO: 10/ 224742 [PALM]  
DATE FILED: August 21, 2002

RELATED-US-APPL-DATA:

Application 10/224742 is a division-of US application 09/122926, filed July 27, 1998, PENDING  
Application is a non-provisional-of-provisional application 60/054012, filed July 29, 1997,

INT-CL: [07] A61 K 38/13, A61 K 31/70, A61 K 31/58, A61 K 31/56, A61 K 31/496, A61 K 31/4745,  
A61 K 31/337, A61 K 31/355

US-CL-PUBLISHED: 424/400; 514/9, 514/176, 514/177, 514/283, 514/449, 514/365, 514/36, 514/458, 514/254.07, 514/389  
US-CL-CURRENT: 424/400; 514/176, 514/177, 514/254.07, 514/283, 514/36, 514/365, 514/389, 514/449, 514/458, 514/9

**ABSTRACT:**

The present invention provides a novel pharmaceutical composition based on the use of a particular oil phase which comprises a lipophilic, pharmaceutically active agent, a mixture of diglyceride and monoglyceride in a ratio of from about 9:1 to about 6:4 by weight (diglyceride:monoglyceride) wherein the diglyceride and monoglyceride are mono- or di-unsaturated fatty acid esters of glycerol having sixteen to twenty-two carbon chain length, one or more pharmaceutically acceptable solvents, and one or more pharmaceutically acceptable surfactants. The composition is in a form of self-emulsifying formulation which provides high concentration and high oral bioavailability for lipophilic compounds.

### CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of the following provisional application: U.S. Serial No. 60/054,078, filed Jul. 29, 1997, under 35 USC 119(e)(i).



DOCUMENT-IDENTIFIER: US 20030044434 A1

TITLE: Self-emulsifying formulation for lipophilic compounds

CLAIMS:

19. The pharmaceutical composition of claim 1 wherein the pharmaceutically acceptable surfactant is Polyoxyl 40 hydrogenated castor oil, Polyoxyl 35 castor oil, Solutol HS-15, Tagat TO, Peglicol 6-oleate, Polyoxyethylene stearates, Poloxamers, Polysorbates, or Saturated Polyglycolyzed Glycerides.



US 20030044434A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0044434 A1****Gao et al.**(43) **Pub. Date: Mar. 6, 2003**(54) **SELF-EMULSIFYING FORMULATION FOR  
LIPOPHILIC COMPOUNDS****Publication Classification**(76) **Inventors: Ping Gao, Portage, MI (US); Walter  
Morozowich, Kalamazoo, MI (US)**

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 38/13; A61K 31/70;**  
**A61K 31/58; A61K 31/56;**  
**A61K 31/496; A61K 31/4745;**  
**A61K 31/337; A61K 31/355**  
(52) **U.S. Cl.** ..... **424/400; 514/9; 514/176;**  
**514/177; 514/283; 514/449;**  
**514/365; 514/36; 514/458;**  
**514/254.07; 514/389**

**Correspondence Address:**  
**Austin W. Zhang**  
**Pharmacia & Upjohn Company**  
**Global Intellectual Property**  
**301 Henrietta Street**  
**Kalamazoo, MI 49001 (US)**

(21) **Appl. No.: 10/224,742**(22) **Filed: Aug. 21, 2002****Related U.S. Application Data**(62) **Division of application No. 09/122,926, filed on Jul.  
27, 1998.**(60) **Provisional application No. 60/054,012, filed on Jul.  
29, 1997.**(57) **ABSTRACT**

The present invention provides a novel pharmaceutical composition based on the use of a particular oil phase which comprises a lipophilic, pharmaceutically active agent, a mixture of diglyceride and monoglyceride in a ratio of from about 9:1 to about 6:4 by weight (diglyceride:monoglyceride) wherein the diglyceride and monoglyceride are mono- or di-unsaturated fatty acid esters of glycerol having sixteen to twenty-two carbon chain length, one or more pharmaceutically acceptable solvents, and one or more pharmaceutically acceptable surfactants. The composition is in a form of self-emulsifying formulation which provides high concentration and high oral bioavailability for lipophilic compounds.

DOCUMENT-IDENTIFIER: US 6531139 B1

TITLE: Self-emulsifying formulation for lipophilic compounds

## CLAIMS:

1. A pharmaceutical composition comprising; (a) a lipophilic, pharmaceutically active agent, (b) a mixture consisting essentially of diglyceride and monoglyceride in a ratio of from about 9:1 to about 6:4 by weight (diglyceride:monoglyceride) wherein the diglyceride and monoglyceride are mono- or di-unsaturated fatty acid esters of glycerol having sixteen to twenty-two carbon chain length, (c) one or more pharmaceutically acceptable solvents, and (d) one or more pharmaceutically acceptable surfactants; wherein the solvent is propylene glycol, polypropylene glycol, polyethylene glycol, glycerol, ethanol, dimethyl isosorbide, glycofurol, propylene carbonate, dimethyl acetamide, or a mixture thereof; wherein said surfactant is Polyoxyl 40 hydrogenated castor oil, Polyoxyl 35 castor oil, polyethylene glycol 12-hydroxy stearate, polyoxyethylene glyceryl trioleate, Peglicol 6-oleate, Polyoxyethylene stearates, Poloxamers, Polysorbates or Saturated Polyglycolized Glycerides; with the proviso that said lipophilic, pharmaceutically active agent has a LOG P value  $\geq 2$ , an intrinsic aqueous solubility  $\leq 0.1$  in the pH range of 1 to 8 and a solubility greater than 1 mg/ml in the mixture of b, c and d as defined herein.



US006531139B1

(12) **United States Patent**  
**Gao et al.**

(10) **Patent No.:** **US 6,531,139 B1**  
(45) **Date of Patent:** **\*Mar. 11, 2003**

(54) **SELF-EMULSIFYING FORMULATION FOR LIPOPHILIC COMPOUNDS**

(75) **Inventors:** **Ping Gao, Portage; Walter Morozowich, Kalamazoo, both of MI (US)**

(73) **Assignee:** **Pharmacia & Upjohn Company, Kalamazoo, MI (US)**

(\*) **Notice:** This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) **Appl. No.:** **09/122,926**

(22) **Filed:** **Jul. 27, 1998**

#### **Related U.S. Application Data**

(60) **Provisional application No. 60/054,078, filed on Jul. 29, 1997.**

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 9/48; A61K 31/44; A61K 31/35**

(52) **U.S. Cl.** ..... **424/400; 424/451; 424/455; 424/456; 514/2; 514/9; 514/169; 514/315; 514/336; 514/456; 514/459; 514/460; 514/557; 514/558; 514/560; 514/573; 514/937; 514/11**

(58) **Field of Search** ..... **424/400, 451, 424/455, 456; 514/2, 9, 169, 459, 460, 557, 558, 560, 937-943, 315, 336, 456, 573**

(56) **References Cited**

#### **U.S. PATENT DOCUMENTS**

4,230,702 A 10/1980 Eckert et al.  
5,504,068 A \* 4/1996 Komiya ..... 514/11  
5,639,724 A \* 6/1997 Cavanak ..... 514/11  
5,645,856 A \* 7/1997 Lacy ..... 424/455

#### **FOREIGN PATENT DOCUMENTS**

EP 0 267 617 A1 5/1988  
GB 2 222 770 A 3/1990  
GB 2 228 198 A 8/1990  
GB 2 257 359 B 10/1996  
WO 94/08603 \* 4/1994  
WO WO95/30670 11/1995  
WO WO96/39142 12/1996  
WO WO98/22106 5/1998

\* cited by examiner

**Primary Examiner**—Gollamudi S. Kishore

(74) **Attorney, Agent, or Firm**—Lucy X. Yang; Austin W. Zhang; Sidney B. Williams, Jr.

(57) **ABSTRACT**

The present invention provides a novel pharmaceutical composition based on the use of a particular oil phase which comprises a lipophilic, pharmaceutically active agent, a mixture of diglyceride and monoglyceride in a ratio of from about 9:1 to about 6:4 by weight (diglyceride:monoglyceride) wherein the diglyceride and monoglyceride are mono- or di-unsaturated fatty acid esters of glycerol having sixteen to twenty-two carbon chain length, one or more pharmaceutically acceptable solvents, and one or more pharmaceutically acceptable surfactants. The composition is in a form of self-emulsifying formulation which provides high concentration and high oral bio-availability for lipophilic compounds.

**39 Claims, No Drawings**

First Hit   Fwd Refs

L8: Entry 99 of 160

File: USPT

Sep 19, 2000

DOCUMENT-IDENTIFIER: US 6121313 A

TITLE: Pharmaceutical composition in a form of self-emulsifying formulation for lipophilic compounds

## CLAIMS:

19. The pharmaceutical composition of claim 1 wherein the pharmaceutically acceptable surfactant is Polyoxyl 40 hydrogenated castor oil, Polyoxyl 35 castor oil, Solutol HS-15, Tagat TO, Peglicol 6-oleate, Polyoxyethylene stearates, Poloxamers, Polysorbates, or Saturated Polyglycolized Glycerides.



US006121313A

**United States Patent** [19]

Gao et al.

[11] **Patent Number:** **6,121,313**[45] **Date of Patent:** **Sep. 19, 2000**

[54] **PHARMACEUTICAL COMPOSITION IN A FORM OF SELF-EMULSIFYING FORMULATION FOR LIPOPHILIC COMPOUNDS**

[75] **Inventors:** Ping Gao, Portage; Walter Morozowich, Kalamazoo, both of Mich.

[73] **Assignee:** Pharmacia & Upjohn Company, Kalamazoo, Mich.

[21] **Appl. No.:** 09/123,069

[22] **Filed:** Jul. 27, 1998

**Related U.S. Application Data**

[60] **Provisional application No.** 60/054,078, Jul. 29, 1997.

[51] **Int. Cl.<sup>7</sup>** ..... **A61K 31/35; A61K 31/19; A61K 31/20**

[52] **U.S. Cl.** ..... **514/459; 514/460; 514/557; 514/558; 514/560**

[58] **Field of Search** ..... **514/459, 460, 514/557, 558, 560**

[56] **References Cited**

**U.S. PATENT DOCUMENTS**

4,230,702 10/1980 Eckert et al. .... 424/242

**FOREIGN PATENT DOCUMENTS**

0 267 617 A1	5/1988	European Pat. Off. ....	A61K 47/00
0267617	5/1988	European Pat. Off. ....	A61K 47/00
2 222 770	3/1990	United Kingdom ....	A61K 9/10
2 228 198	8/1990	United Kingdom ....	A61K 37/02
2 257 359	10/1996	United Kingdom ....	A61K 38/13
WO95/30670	11/1995	WIPO .....	C07D 309/32
WO96/39142	12/1996	WIPO .....	A61K 31/47
WO98/22106	5/1998	WIPO .....	A61K 31/425

*Primary Examiner*—Russell Travers

*Attorney, Agent, or Firm*—Lucy X. Yang

[57] **ABSTRACT**

The present invention provides a novel pharmaceutical composition based on the use of a particular oil phase which comprises a pyranone compound as a pharmaceutically active agent, a mixture of diglyceride and monoglyceride in a ratio of from about 9:1 to about 6:4 (diglyceride:monoglyceride) wherein the diglyceride and monoglyceride are mono- or di-unsaturated fatty acid esters of glycerol having sixteen to twenty-two carbon chain length, one or more pharmaceutically acceptable solvents, and one or more pharmaceutically acceptable surfactants. The composition is in a form of a self-emulsifying formulation which provides high concentration and high oral bioavailability for lipophilic pyranone compounds.

**34 Claims, No Drawings**

US-PAT-NO: 5681552

DOCUMENT-IDENTIFIER: US 5681552 A

TITLE: Clear cosmetic stick composition containing a combination of anionic and non-ionic surfactants

DATE-ISSUED: October 28, 1997

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Shevade; Makarand	Plainsboro	NJ		
Moghe; Bhalchandra D.	Scotch Plains	NJ		
Kasat; Radhakrishna B.	Bellemead	NJ		

US-CL-CURRENT: 424/65, 424/78.02, 424/78.08, 424/78.18, 512/1, 514/944, 514/946

## CLAIMS:

We claim:

1. A clear stick composition, comprising alcohol and water, and a soap gelling agent, the soap gelling agent being included in an amount so as to gel and to form the stick composition, the composition further including, as a clarifying agent, a combination of both (a) at least one anionic surface active agent and (b) at least one non-ionic surface active agent, the at least one non-ionic surface active agent including a straight chain primary alkoxyated alcohol, other than nonoxynol-10, cetareth-12, cetareth-20, cetareth-30, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, PPG-2 cetareth-9, oleth-10-, oleth-5 and PEG-40 castor oil, the combination of anionic and non-ionic surface active agents being included in the composition in an amount so as to provide a clear stick composition, the at least one anionic surface active agent being included in the composition in an amount of 2-8% by weight, of the total weight of the composition, and the at least one non-ionic surface active agent being included in the composition in an amount of 2-8% by weight, of the total weight of the composition.

2. A clear stick composition according to claim 1, wherein the composition further includes a deodorant effective amount of a deodorant active material, whereby the composition is a clear deodorant stick composition.

3. A clear stick composition according to claim 2, wherein said deodorant active material is at least one selected from the group consisting of fragrances and antibacterial agents.

4. A clear stick composition according to claim 1, wherein the alcohol includes propylene glycol, and the soap gelling agent includes alkali metal salts of fatty acids.

5. A clear stick composition according to claim 4, wherein the composition includes, in percent by weight of the total weight of the composition, 55-80% propylene glycol, 9-25% water and 4-10% soap gelling agent.

6. A clear stick composition according to claim 1, wherein the composition includes, in percent by weight of the total weight of the composition, 4-6% of the at least one anionic surface active agent and 2-4% of the at least one

non-ionic surface active agent.

7. A clear stick composition according to claim 1, wherein said straight chain primary alkoxyated alcohol is a straight chain primary ethoxylated alcohol.

8. A clear stick composition according to claim 1, wherein said at least one non-ionic surface active agent has a hydrophile-lipophile balance of at least 24.

9. A clear stick composition according to claim 8, wherein the straight chain primary alkoxyated alcohol is a straight chain primary ethoxylated alcohol.

10. A clear stick composition according to claim 8, wherein the at least one anionic surface active agent includes sodium laureth-13 carboxylate.

11. A clear stick composition according to claim 10, wherein the at least one non-ionic surface active agent is selected from the group consisting of Ceteareth-55, C14-15 Pareth-2.25 and C14-15 Pareth-13.

12. A clear stick composition according to claim 11, wherein the at least one non-ionic surface active agent includes Ceteareth-55.

13. A clear stick composition according to claim 7, wherein the composition further includes a deodorant effective amount of a deodorant active material, whereby the composition is a clear deodorant stick composition.

14. A clear deodorant stick composition according to claim 13, wherein said deodorant active material is at least one selected from the group consisting of fragrances and antibacterial agents.

15. A clear stick composition according to claim 1, wherein the composition further includes a deodorant effective amount of a deodorant active material, whereby the composition is a clear deodorant stick composition.

16. A clear deodorant stick composition according to claim 15, wherein said deodorant active material is at least one selected from the group consisting of fragrances and antibacterial agents.

17. A clear stick composition according to claim 1, wherein the at least one anionic surface active agent is selected from the group consisting of long-chain fatty acids, sulfosuccinates, alkyl sulfates, phosphates and sulfonates.

18. A clear stick composition according to claim 1, wherein the at least one non-ionic surface active agent includes polyethylene oxide groups.

19. A clear stick composition according to claim 1, wherein the soap gelling agent includes salts of fatty acids, the fatty acids having a carbon chain length in a range of C.sub.12 -C.sub.22, at least some of the salts being salts of fatty acids having carbon chain length of at least one of C.sub.20 and C.sub.22.

20. A clear stick composition according to claim 1, wherein the at least one anionic surface active agent includes a sodium salt of methyl carboxy derivatives of ethoxylated lauryl alcohol.

21. A clear stick composition according to claim 1, wherein the soap gelling



agent includes sodium laurate, sodium myristate, sodium palmitate, sodium stearate, sodium arachidate and sodium behenate.

22. A packaged clear deodorant stick composition, comprising the clear deodorant stick composition of claim 15 in a dispensing package, wherein the dispensing package is made of a styrene-butadiene copolymer.

23. A packaged clear deodorant stick composition according to claim 22, wherein the dispensing package is clear.

24. A packaged clear stick composition, comprising the clear stick composition of claim 1 in a dispensing package, wherein the dispensing package is made of a styrene-butadiene copolymer.

25. A packaged clear deodorant stick composition according to claim 24, wherein the dispensing package is clear.

26. A method of reducing body malodor, comprising rubbing the clear deodorant stick composition of claim 15 on axillary regions of a human body.

27. A method of reducing body malodor, comprising rubbing the clear deodorant stick composition of claim 13 on axillary regions of human body.

28. A method of reducing body malodor, comprising rubbing the clear deodorant stick composition of claim 2 on axillary regions of a human body.

29. A clear stick composition, comprising alcohol and water, and a soap gelling agent, the soap gelling agent being included in an amount so as to gel and to form the stick composition, the composition further including, as a clarifying agent, a combination of both (a) at least one anionic surface active agent and (b) at least one non-ionic surface active agent, the at least one non-ionic surface active agent including an alkoxylate homopolymer, other than nonoxynol-10, cetareth-12, cetareth-20, cetareth-30, PEG-5 cocamide, PEG-40 hydrogenated castor oil, PEG-60 hydrogenated castor oil, oleth-5, oleth-10, PEG-40 castor oil, polysorbate 20, polysorbate 60, polysorbate 80 and PEG-8 stearate, the combination of anionic and non-ionic surface active agents being included in the composition in an amount so as to provide a clear stick composition, the at least one anionic surface active agent being included in the composition in an amount of 2-8% by weight, of the total weight of the composition, and the at least one non-ionic surface active agent being included in the composition in an amount of 2-8% by weight, of the total weight of the composition.

30. A clear stick composition according to claim 29, wherein the composition includes, in percent by weight of the total weight of the composition, 4-6% of the at least one anionic surface active agent and 2-4% of the at least one non-ionic surface active agent.

31. A clear stick composition according to claim 29, wherein said alkoxylate homopolymer is an ethoxylate homopolymer.

32. A clear stick composition according to claim 29, wherein said at least one non-ionic surface active agent has a hydrophile-lipophile balance of at least 24.

33. A clear stick composition, comprising alcohol and water, and a soap gelling agent, the soap gelling agent being included in an amount so as to gel

and to form the stick composition, the composition further including, as a clarifying agent, a combination of both (a) at least one anionic surface active agent and (b) at least one non-ionic surface active agent, the at least one non-ionic surface active agent having a hydrophile-lipophile balance of at least 24, the combination of anionic and non-ionic surface active agents being included in the composition in an amount so as to provide a clear stick composition.

34. A clear stick composition according to claim 33, wherein the at least one anionic surface active agent and the at least one non-ionic surface active agent are each included in the composition in an amount of 2-8% by weight, of the total weight of the composition.

35. A clear stick composition according to claim 34, wherein the composition includes, in percent by weight of the total weight of the composition, 4-6% of the at least one anionic surface active agent and 2-4% of the at least one non-ionic surface active agent.



US005681552A

**United States Patent** [19]

Shevade et al.

[11] Patent Number: 5,681,552

[45] Date of Patent: Oct. 28, 1997

[54] **CLEAR COSMETIC STICK COMPOSITION CONTAINING A COMBINATION OF ANIONIC AND NON-IONIC SURFACTANTS**

[75] Inventors: Makarand Shevade, Plainsboro;  
Bhalchandra D. Moghe, Scotch Plains;  
Radhakrishna B. Kasat, Bellemcad, all  
of NJ.

[73] Assignee: The Mennen Company, Morristown,  
NJ.

[21] Appl. No.: 448,104

[22] Filed: May 23, 1995

[51] Int. Cl.<sup>6</sup> A61K 7/32

[52] U.S. Cl. 424/65; 424/78.02; 424/78.08;  
424/78.18; 512/1; 514/944; 514/946

[58] Field of Search 424/65, 78.02,  
424/78.08, 78.18; 512/1; 514/944, 946

[56] **References Cited****U.S. PATENT DOCUMENTS**

4,122,029 10/1978 Gee et al. .  
4,151,272 4/1979 Gary et al. .  
4,265,878 5/1981 Keil .  
4,268,498 5/1981 Gedeon et al. .  
4,504,465 3/1985 Sampson et al. .  
4,759,924 7/1988 Luebke et al. .  
4,822,602 4/1989 Sabatelli .  
5,114,717 5/1992 Kuznitz et al. .  
5,128,123 7/1992 Brewster et al. . 424/65

5,198,218 3/1993 Kuznitz et al. . 424/401  
5,316,761 5/1994 Brazinsky . 424/65  
5,424,070 6/1995 Kasat et al. . 424/401  
5,458,880 10/1995 Kasat et al. . 424/401  
5,462,736 10/1995 Rech et al. . 424/401  
5,463,098 10/1995 Giovanniello et al. . 424/66

**FOREIGN PATENT DOCUMENTS**

0450597 10/1991 European Pat. Off. .  
062338A2 4/1994 European Pat. Off. .  
WO94/27567 12/1994 WIPO .


Primary Examiner—Terressa Mosley

Attorney, Agent, or Firm—Antonelli, Terry, Stout & Kraus,  
LLP

[57] **ABSTRACT**

Disclosed are clear, soap-gelled cosmetic (e.g., deodorant) stick compositions, containing alcohol and water, and including a combination of anionic (e.g., sodium laureth-13 carboxylate) and non-ionic (e.g., Ceteareth-55) surfactants as a clarifying agent. The non-ionic surfactants preferably include ethoxylated alcohols, and preferably have an HLB $\geq$ 24. The compositions can include various cosmetically active materials, including deodorant active materials (fragrance, Triclosan, etc.). The compositions have superior clarity and maintain superior clarity over extended periods of time, have improved (smoother) glide over the skin and a smoother surface, exhibit reduced syneresis in K-resin and polypropylene dispensing packages, and are easy to manufacture and have high gelling temperatures.

35 Claims, 5 Drawing Sheets



US-PAT-NO: 5189066

DOCUMENT-IDENTIFIER: US 5189066 A

TITLE: Pharmaceutical compositions of tebufelone

DATE-ISSUED: February 23, 1993

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Kelm; Gary R.	Cincinnati	OH		
Bruns; Alan E.	Okeana	OH		

US-CL-CURRENT: 514/678; 514/689, 514/941, 514/960, 514/961

## CLAIMS:

What is claimed is:

1. A composition consisting essentially of 1-3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl-5-hexyn-1-one as a drug active at a concentration of at least about 15% of the composition, and the balance a pharmaceutically-acceptable vehicle, the composition having the following properties:

(1) being a homogeneous liquid at 37.degree. C.,

(2) providing solubilization of the drug active at a level of at least 1 mg/mL in 0.1N HCl at 20.degree. C., and

(3) providing solubilization of 20 mg of the drug active in 500 mL of simulated intestinal fluid in 5 minutes;

the vehicle comprising a surfactant or mixture of surfactants, the vehicle having the following properties:

(a) being a homogeneous liquid at 37.degree. C.,

(b) having an HLB of from about 9 to about 13,

(c) forming a stable dispersion in water at 20.degree. C. at concentrations of 10%;

whereby the absorption of the drug active from the gastrointestinal tract is substantially greater for the composition when perorally administered than for conventional solid dosage forms of the drug active.

2. The composition of claim 1 wherein the vehicle additionally has the following properties:

(d) being soluble in isopropanol at 20.degree. C. at concentrations of 10%, and

(e) being soluble in cottonseed oil at 20.degree. C. at concentrations of 1%.

3. The composition of claim 1 wherein the surfactant is selected from the group consisting of polysorbate 81, poloxamer 182, poloxamer 183, poloxamer 184, PEG-25 glyceryl trioleate, and polyoxyl 35 castor oil, and mixtures thereof.

4. The composition of any of claims 1, 2 or 3 wherein the vehicle consists essentially of the surfactant or surfactant mixture.

5. The composition of claim 1 wherein the vehicle also comprises a lipophilic solvent selected from the group consisting of a triglyceride or a mixture of triglycerides having fatty chains of from about 6 to about 10 carbon atoms, straight-chain saturated fatty acids having from about 6 to about 10 carbon atoms, and straight-chain unsaturated fatty acids having from about 12 to about 18 carbon atoms, and mixtures thereof.

6. The compositions of claim 5 wherein the surfactant is selected from the group consisting of polysorbate 80, polysorbate 81, poloxamer 182, poloxamer 183, poloxamer 184, PEG-25 glyceryl trioleate and polyoxyl 35 castor oil, and mixtures thereof.

7. The composition of claim 6 wherein the vehicle consists essentially of from about 25% to about 85% of a lipophilic solvent which is selected from the group consisting of caprylic/capric triglyceride, oleic acid and linoleic acid; and from about 15% to about 75% of the surfactant.

8. The composition of claim 3 consisting essentially of from about 15% to about 20% tebufelone, from about 35% to about 45% polyoxyl 35 castor oil, and from about 35% to about 45% poloxamer 182.

9. The composition of claim 5 consisting essentially of from about 15% to about 20% tebufelone, from about 4% to about 6% polyoxyl 35 castor oil, from about 20% to about 25% poloxamer 182, and from about 50% to about 60% caprylic/capric triglyceride.

10. A composition consisting essentially of 1-3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl-5-hexyn-1-one as a drug active at a concentration of at least about 15% of the composition, and the balance a pharmaceutically-acceptable vehicle, the composition having the following properties:

(1) being a homogeneous liquid at 37.degree. C.,

(2) providing solubilization of the drug active of at least 1 mg/mL in 0.1N HCl at 20.degree. C., and

(3) providing solubilization of 20 mg of the drug active in 500 mL of simulated intestinal fluid in 5 minutes;

the vehicle comprising triglycerides interesterified with polyethylene glycol, such that its HLB is from about 3 to about 7; whereby the absorption of the drug active from the gastrointestinal tract is substantially greater for the composition when perorally administered than for conventional solid dosage forms of the drug active.

11. The composition of claim 10 wherein the triglycerides interesterified with polyethylene glycol are glycolysed ethoxylated glycerides obtained by partial hydrolysis of natural vegetable oils.

12. The composition of claim 11 wherein the triglycerides interesterified with polyethylene glycol are glycolysed ethoxylated glycerides obtained by partial hydrolysis of corn oil with polyethylene glycol 400.
13. The composition of any of claims 10, 11 or 12 wherein the vehicle consists essentially of the triglycerides interesterified with polyethylene glycol.
14. A pharmaceutical unit dosage form comprising a composition of any of claims 3, 6, 8, 9 or 12 in a soft gelatin capsule shell.
15. A pharmaceutical unit dosage form comprising a composition of any of claims 3, 6, 8, 9 or 12 in a sealed hard gelatin capsule shell.
16. The composition of claim 1 wherein the surfactant is 100% polysorbate 81.
17. The composition of claim 1 wherein the surfactant is 100% PEG-25 glyceryl trioleate.
18. The composition of claim 1 wherein the surfactant is 100% poloxamer 183.
19. The composition of claim 1 wherein the surfactant is 100% poloxamer 184.
20. The composition of claim 1 wherein the surfactant is from about 25% to about 75% poloxyl 35 castor oil and from about 25% to about 75% poloxamer 182.
21. The composition of claim 1 wherein the surfactant is from about 10% to about 25% polysorbate 80 and from about 75% to about 90% poloxamer 182.
22. A method for obtaining good absorption of the drug active from the gastrointestinal tract by perorally administering the composition of any of claims 1, 3, 6, 10 or 11.



US005189066A

**United States Patent** [19]

Kelm et al.

[11] Patent Number: **5,189,066**[45] Date of Patent: **Feb. 23, 1993**[54] **PHARMACEUTICAL COMPOSITIONS OF  
TEBUFELONE**[75] Inventors: **Gary R. Kelm, Cincinnati; Alan E.  
Bruns, Okeana, both of Ohio**[73] Assignee: **The Procter & Gamble Company,  
Cincinnati, Ohio**[21] Appl. No.: **732,951**[22] Filed: **Jul. 19, 1991****Related U.S. Application Data**

[63] Continuation of Ser. No. 440,178, Nov. 22, 1989, abandoned.

[51] Int. Cl.<sup>5</sup> ..... **A61K 31/12**[52] U.S. Cl. .... **514/678; 514/689;  
514/960; 514/961; 514/941**[58] Field of Search ..... **514/678, 689, 960, 961,  
514/941**[56] **References Cited****U.S. PATENT DOCUMENTS**

2,988,484	6/1961	Barsky et al. ....	424/107
3,867,521	2/1975	Miskel et al. ....	424/37
4,156,219	5/1979	Sezaki .....	424/118
4,325,942	4/1982	Taki et al. ....	424/94.1
4,690,816	9/1987	Hata et al. ....	424/456
4,690,823	9/1987	Lohner et al. ....	424/456
4,708,966	11/1987	Loomans et al. ....	514/689
4,786,495	11/1988	Bird et al. ....	424/81
4,797,288	1/1989	Sharma et al. ....	424/476
4,827,062	5/1989	Sacki et al. ....	514/690

4,847,303	7/1989	Loomans et al. ....	549/689
4,849,428	7/1989	Dobson et al. ....	549/307
4,853,379	8/1989	Shroot et al. ....	514/179

**OTHER PUBLICATIONS**

Griffin, W. C., "Classification of Surface-Active Agents by 'HLB'", *Journal of the Society of Cosmetic Chemists*, vol. 1, No. 5 (1949), pp. 311-326.  
 CTFA Cosmetic Ingredient Dictionary, Third Edition (1984), N. F. Estrin, P. A. Crosely and C. R. Haynes, Eds., The Cosmetic Toiletory and Fragrance Association, Inc., Washington, D.C., pp. 39-40, 211, 238 & 247.  
 The National Formulary, 17th Edition (1990), The United States Pharmacopeial Convention, Inc., Rockville, Md., p. 1966.

*Primary Examiner*—Frederick E. Waddell*Assistant Examiner*—T. J. Criares*Attorney, Agent, or Firm*—M. B. Graff, IV; J. D. Schaeffer; T. H. O'Flaherty[57] **ABSTRACT**

The subject invention involves compositions, consisting essentially of the drug active 1-3,5-bis(1,1-dimethyl-ethyl)-4-hydroxyphenyl-5-hexyn-1-one (tebufelone) at a concentration of at least about 15%, and the balance a pharmaceutically-acceptable vehicle. The vehicle is formulated such that the compositions are homogeneous liquids at 37° C. and provide good solubilization of the drug active.

**22 Claims, No Drawings**